

FREE BROWNIAN MOTION AND FREE CONVOLUTION SEMIGROUPS: MULTIPLICATIVE CASE

PING ZHONG

ABSTRACT. We consider a pair of probability measures μ, ν on the unit circle such that $\Sigma_\lambda(\eta_\nu(z)) = z/\eta_\mu(z)$. We prove that the same type of equation holds for any $t \geq 0$ when we replace ν by $\nu \boxtimes \lambda_t$ and μ by $\mathbb{M}_t(\mu)$, where λ_t is the free multiplicative analogue of the normal distribution on the unit circle of $\mathbb{C}$ and $\mathbb{M}_t$ is the map defined by Arizmendi and Hasebe. These equations are a multiplicative analogue of equations studied by Belinschi and Nica. In order to achieve this result, we study infinite divisibility of the measures associated with subordination functions in multiplicative free Brownian motion and multiplicative free convolution semigroups. We use the modified $\mathcal{S}$ -transform introduced by Raj Rao and Speicher to deal with the case that ν has mean zero. The same type of the result holds for convolutions on the positive real line. We also obtain some regularity properties for the free multiplicative analogue of the normal distributions.

1. INTRODUCTION

Let $\mathcal{M}_{\mathbb{R}}$ be the set of probability measures on $\mathbb{R}$. For every $t \geq 0$, Belinschi and Nica [11] defined a family of maps $\mathbb{B}_t : \mathcal{M}_{\mathbb{R}} \rightarrow \mathcal{M}_{\mathbb{R}}$ by setting

$$\mathbb{B}_t(\mu) = (\mu^{\boxplus(t+1)})^{\boxplus \frac{1}{t+1}}, \quad \mu \in \mathcal{M}_{\mathbb{R}}.$$

These maps have several remarkable properties. For any $t \geq 0$, $\mathbb{B}_t$ is an endomorphism of $(\mathcal{M}_{\mathbb{R}^+}, \boxtimes)$, where $\mathcal{M}_{\mathbb{R}^+}$ is the set of probability measures on $[0, +\infty)$ and $\boxtimes$ is free multiplicative convolution. $\{\mathbb{B}_t\}_{t \geq 0}$ is a semigroup and $\mathbb{B}_1$ is the Boolean to free Bercovici-Pata bijective map.

The maps $\mathbb{B}_t$ have strong connections with $\boxplus$ -infinite divisibility. They are also connected to free Brownian motion and additive free convolution semigroups. For $\mu \in \mathcal{M}_{\mathbb{R}}$, we denote by G_μ the Cauchy transform of μ and by F_μ the reciprocal Cauchy transform of μ . Given a pair of probability measures $\mu, \nu \in \mathcal{M}_{\mathbb{R}}$ such that

$$G_\nu(z) = z - F_\mu(z), \quad z \in \mathbb{C}^+,$$

then we have

$$(1.1) \quad G_{\nu \boxplus \gamma_t}(z) = z - F_{\mathbb{B}_t(\mu)}(z), \quad t > 0, \quad z \in \mathbb{C}^+,$$

where γ_t is the semi-circular distribution with variance t . This result was generalized to the context of multi-variable case in [10], [12] and [26]. An equivalent form of (1.1) was used to prove the superconvergence theorem in [32]. In a series of papers

[1], [2], [3] and [4], Anshelevich generalized the above correspondence of $\mu \leftrightarrow \nu$ and $\mathbb{B}_t(\mu) \leftrightarrow \nu \boxtimes \gamma_t$ to the context of two-state probability spaces. Motivated by these generalizations and applications, we study in this article the analogue of these equations for multiplicative free convolution.

Throughout this article, we denote by $\mathbb{T}$ the unit circle of $\mathbb{C}$, by $\mathcal{M}_{\mathbb{T}}$ the set of probability measures on $\mathbb{T}$, and by $\mathcal{M}_*$ the set of probability measures on $\mathbb{C}$ with nonzero mean. We also set

$$\mathcal{M}_{\mathbb{T}}^* = \{\mu \in \mathcal{M}_{\mathbb{T}} \cap \mathcal{M}_* : \eta_{\mu}(z) \neq 0, \forall z \in \mathbb{D} \setminus \{0\}\}.$$

It was shown in [8] that one can define multiplicative free convolution power $\mu^{\boxtimes t}$ for $\mu \in \mathcal{M}_{\mathbb{T}}^*$ and $t > 1$.

In [6], a family of maps $\mathbb{M}_t$, which is the analogue of the semigroup $\mathbb{B}_t$, was defined for the probability measures in $\mathcal{M}_{\mathbb{T}}^*$. The definition of $\mathbb{M}_t$ in [6] is more general, and we only need a simpler form defined as follows. Given $\mu \in \mathcal{M}_{\mathbb{T}}^*$ which has positive mean, then for $t \geq 0$, the map $\mathbb{M}_t$ is defined by

$$\mathbb{M}_t(\mu) = (\mu^{\boxtimes(t+1)})^{\boxtimes \frac{1}{t+1}},$$

where the convolution power $\mu^{\boxtimes(t+1)}$ and the measure $\mathbb{M}_t(\mu)$ are chosen in a way such that they have positive means.

We then state one of our main theorems.

Theorem 1.1. *Given a pair of probability measures $\mu \in \mathcal{M}_{\mathbb{T}}^*$ and $\nu \in \mathcal{M}_{\mathbb{T}}$ such that*

$$(1.2) \quad \Sigma_{\lambda}(\eta_{\nu}(z)) = \frac{z}{\eta_{\mu}(z)}, z \in \mathbb{D}.$$

Then we have

$$(1.3) \quad \Sigma_{\lambda}(\eta_{\nu \boxtimes \lambda_t}(z)) = \frac{z}{\eta_{\mathbb{M}_t(\mu)}(z)}, z \in \mathbb{D},$$

where λ_t is the analogue of the normal distribution on $\mathbb{T}$ with $\Sigma_{\lambda_t}(z) = \exp(\frac{t}{2} \frac{1+z}{1-z})$ and $\lambda = \lambda_1$.

In order to prove Theorem 1.1, we consider two semigroups $\nu \boxtimes \lambda_t$ and $\mu^{\boxtimes(t+1)}$ for all $t \geq 0$. It is well-known that $\eta_{\nu \boxtimes \lambda_t}$ and $\eta_{\mu^{\boxtimes(t+1)}}$ are subordinated to η_{ν} and η_{μ} respectively. We prove that the subordination functions are η -transforms of some $\boxtimes$ -infinitely divisible probability measures on $\mathbb{T}$. It turns out that the equation $\Sigma_{\lambda}(\eta_{\nu}(z)) = z/\eta_{\mu}(z)$ means that the subordination function of $\eta_{\nu \boxtimes \lambda_t}$ with respect to η_{ν} and the subordination function of $\eta_{\mu^{\boxtimes(t+1)}}$ with respect to η_{μ} are the same. The proof of Theorem 1.1 will be given in Subsection 3.5.

Given $\mu \in \mathcal{M}_{\mathbb{T}}$, we prove that if $\mu^{\boxtimes t}$ can be defined and $\eta_{\mu^{\boxtimes t}}$ is subordinated to η_{μ} for all $t > 1$, then $\mu \in \mathcal{M}_{\mathbb{T}}^*$; in addition, we prove that for nontrivial measures $\mu \in \mathcal{M}_{\mathbb{T}}$ and $\nu \in \mathcal{ID}(\boxtimes, \mathbb{T})$, the density functions of the measures $\mu \boxtimes \nu_t$ and $\mu^{\boxtimes t}$ converge to $1/2\pi$ uniformly as $t \rightarrow \infty$.

To deal with the case that $\nu \in \mathcal{M}_T \setminus \mathcal{M}_*$, we use the modified $\mathcal{S}$ -transform [5, 28] to study subordination functions. In this case, the subordination function of $\eta_{\nu \boxtimes \lambda_t}$ with respect to η_ν is generally not unique. However, we can prove that there exists a unique subordination function satisfying certain properties (see Theorem 3.11), which implies that $\Sigma_{\rho_t}(z) = \Sigma_{\lambda_t}(\eta_\nu(z))$, where ρ_t is the measure associated with the subordination function of $\eta_{\nu \boxtimes \lambda_t}$ with respect to η_ν .

Similar results to Theorem 1.1 for multiplicative convolution on $\mathcal{M}_{\mathbb{R}^+}$ are also valid. The proof for this case is much simpler because of the uniqueness of multiplicative convolution powers and the uniqueness of subordination functions.

Finally, we describe the density functions of the free multiplicative analogue of the normal distributions. For λ_t ($t > 0$) the free multiplicative analogue of the normal distributions on $\mathbb{T}$, we prove that λ_t is unimodal.

This article is organized as follows. After this introductory section, we describe some backgrounds in the additive case in Section 2. In Section 3, we consider multiplicative free and multiplicative Boolean convolution on $\mathcal{M}_T$, and prove our main theorems. Section 4 is devoted to studying multiplicative free and multiplicative Boolean convolution on $\mathcal{M}_{\mathbb{R}^+}$. The regularity properties of the free multiplicative analogue of the normal distributions are discussed in Section 5.

2. BACKGROUND: ADDITIVE CASE

2.1. Additive free convolution and additive Boolean convolution. For a measure $\mu \in \mathcal{M}_{\mathbb{R}}$, we define the Cauchy transform $G_\mu : \mathbb{C}^+ \rightarrow \mathbb{C}^-$ by

$$G_\mu(z) = \int_{-\infty}^{+\infty} \frac{1}{z-t} d\mu(t), \quad z \in \mathbb{C}^+.$$

We set $F_\mu(z) = 1/G_\mu(z)$, $z \in \mathbb{C}^+$, so that $F_\mu : \mathbb{C}^+ \rightarrow \mathbb{C}^+$ is analytic.

The following result in [16] characterizes those functions which are reciprocal Cauchy transforms of probability measures.

Proposition 2.1. *Let $F : \mathbb{C}^+ \rightarrow \mathbb{C}^+$ be an analytic function. Then the following assertions are equivalent.*

- (1) *There exists a probability measure μ on $\mathbb{R}$ such that $F(z) = F_\mu(z)$ in $\mathbb{C}^+$.*
- (2) *There exists $a \in \mathbb{R}$, and a finite positive measure ρ on $\mathbb{R}$ such that*

$$F(z) = a + z + \int_{-\infty}^{+\infty} \frac{1+tz}{t-z} d\rho(t)$$

for all $z \in \mathbb{C}^+$.

- (3) *We have that $\lim_{y \rightarrow +\infty} F(iy)/iy = 1$.*

It was proved in [16] that F_μ is invertible in some domain. More precisely, for two positive numbers M and N , we set

$$\Gamma_{M,N} = \{z \in \mathbb{C}^+ : |x| < My, y > N\}.$$

Then for any $M > 0$, there exists $N > 0$ such that the left inverse F_μ^{-1} of F_μ is defined in $\Gamma_{M,N}$, and then we can define the Voiculescu transform of μ by

$$\varphi_\mu(z) = F_\mu^{-1}(z) - z,$$

for $z \in \Gamma_{M,N}$. For any two measures $\mu, \nu \in \mathcal{M}_\mathbb{R}$, then we have that

$$(2.1) \quad \varphi_{\mu \boxplus \nu}(z) = \varphi_\mu(z) + \varphi_\nu(z)$$

holds in any truncated cone $\Gamma_{M,N}$ where φ_μ, φ_ν and $\varphi_{\mu \boxplus \nu}$ are defined. This remarkable result was discovered by Voiculescu [30] for compactly supported measures and then extended to general cases in [16] [25].

Given $\nu \in \mathcal{M}_\mathbb{R}$, we say that ν is $\boxplus$ -infinitely divisible if for every positive integer n , there exists a probability measure $\nu_{1/n} \in \mathcal{M}_\mathbb{R}$ such that

$$\nu = \underbrace{\nu_{1/n} \boxplus \nu_{1/n} \boxplus \cdots \boxplus \nu_{1/n}}_{n \text{ times}}.$$

It is known [16, 25, 30] that a probability measure ν on $\mathbb{R}$ is $\boxplus$ -infinitely divisible if and only if its Voiculescu transform φ_ν has an analytic extension defined on $\mathbb{C}^+$ with values in $\mathbb{C}^- \cup \mathbb{R}$. We denote by $\mathcal{ID}(\boxplus, \mathbb{R})$ the set of all $\boxplus$ -infinitely divisible probability measures on the real line. If $\nu \in \mathcal{ID}(\boxplus, \mathbb{R})$, then for every $t > 0$, there exists a probability measure ν_t such that $\varphi_{\nu_t}(z) = t\varphi_\nu(z)$ for z in the common domain of φ_ν and φ_{ν_t} .

We would like to mention the following fact.

Proposition 2.2. *If ν is $\boxplus$ -infinitely divisible, let $H(z) = z + \varphi_\nu(z)$, then*

$$(2.2) \quad H(F_\nu(z)) = z$$

holds for $z \in \mathbb{C}^+$. The set $U := \{z \in \mathbb{C}^+ : \Im H(z) > 0\}$ is a simply connected domain with boundary which is a simple curve and H maps $\mathbb{C}^+$ conformally onto U . Moreover, the boundary ∂U is the graph of a function and the function H is continuous up to the real axis.

Proof. The first part of the assertion appears in [16] and [30], and the second part of the assertion follows from the fact that H satisfies the conditions of Proposition 4.7 in [8]. The last part of the assertion is due to Lemma 3.3 in [21] and Proposition 4.7 in [8]. $\square$

Additive Boolean convolution was introduced in [29]. For $\mu \in \mathcal{M}_\mathbb{R}$, we set $E_\mu(z) = z - F_\mu(z)$. For $\mu, \nu \in \mathcal{M}_\mathbb{R}$, the additive Boolean convolution $\mu \uplus \nu$ is characterized by the identity

$$E_\rho(z) = E_\mu(z) + E_\nu(z), \text{ for } z \in \mathbb{C}^+.$$

We can also consider the infinite divisibility with respect to additive Boolean convolution. It turns out that every $\mu \in \mathcal{M}_\mathbb{R}$ is $\uplus$ -infinitely divisible (see [29]). We denote by $\mathcal{ID}(\uplus, \mathbb{R})$ the set of all $\uplus$ -infinitely divisible probability measures on the real line.

2.2. Infinite divisibility and subordination functions. Given $\mu, \nu \in \mathcal{M}_{\mathbb{R}}$, it is known that $F_{\mu \boxplus \nu}$ is subordinated to F_{μ} and F_{ν} , and by Proposition 2.1, we can also regard these subordination functions as the reciprocal Cauchy transforms of probability measures on $\mathbb{R}$.

Definition 2.3. For $\mu, \nu \in \mathcal{M}_{\mathbb{R}}$, the subordination distribution [4, 24, 26] $\mu \boxminus \nu$ (resp. $\nu \boxminus \mu$) is defined to be the unique probability measure in $\mathcal{M}_{\mathbb{R}}$ such that $F_{\mu \boxplus \nu}(z) = F_{\nu}(F_{\mu} \boxminus \nu(z))$ (resp. $F_{\mu \boxplus \nu}(z) = F_{\mu}(F_{\nu} \boxminus \mu(z))$).

Many subordination distributions in semigroups related to free convolution are infinitely divisible (see [4, 26]).

Proposition 2.4. Let $\mu, \nu \in \mathcal{M}_{\mathbb{R}}$.

- (1) We have that $\varphi_{\mu \boxminus \nu}(z) = (\varphi_{\mu} \circ F_{\nu})(z)$.
- (2) If $\mu \in \mathcal{ID}(\boxplus, \mathbb{R})$, then $\mu \boxminus \nu \in \mathcal{ID}(\boxplus, \mathbb{R})$. In particular, $\gamma_t \boxminus \nu \in \mathcal{ID}(\boxplus, \mathbb{R})$ and $\varphi_{\gamma_t \boxminus \nu}(z) = tG_{\nu}(z)$, where γ_t is the semi-circular distribution with variance t .
- (3) If $\nu = \mu \boxplus \nu'$ for $\nu' \in \mathcal{M}_{\mathbb{R}}$, then $\mu \boxminus \nu \in \mathcal{ID}(\boxplus, \mathbb{R})$. In particular, $\mu \boxminus \mu \in \mathcal{ID}(\boxplus, \mathbb{R})$, and $\varphi_{\mu \boxminus \mu}(z) = z - F_{\mu}(z)$.

Proof. Part (1) is Lemma 1 in [4]. Note that $\varphi_{\gamma_t}(z) = t/z$ and $(\varphi_{\mu} \circ F_{\mu})(z) = z - F_{\mu}(z)$. Part (2) and (3) follow from Part 1 and Lemma 2 in [4], see also Corollary 2.3 in [20]. $\square$

The following result was inspired by a question in [4]. I am grateful to Michael Anshelevich for sending me a updated version of the paper [4].

Lemma 2.5. Given $\tau, \rho \in \mathcal{M}_{\mathbb{R}}$, if $\tau \boxminus \rho \in \mathcal{ID}(\boxplus, \mathbb{R})$, then $\rho \boxplus \tau^{\boxplus t}$ is defined for all $t \geq 0$ in the sense that $\varphi_{\rho} + t\varphi_{\tau}$ is the Voiculescu transform of a positive measure. Moreover, we have that $F_{\rho \boxplus (\tau^{\boxplus t})} = F_{\rho} \left(F_{\tau \boxminus \rho}^{\boxplus t}(z) \right)$.

Proof. Let $\sigma = \tau \boxminus \rho$, and $\sigma_t = \sigma^{\boxplus t}$. By Proposition 2.1, there exists a unique probability measure $\mu_t \in \mathcal{M}_{\mathbb{R}}$, such that

$$F_{\mu_t} = F_{\rho} (F_{\sigma_t}(z)).$$

We claim that $\varphi_{\mu_t}(z) = \varphi_{\rho}(z) + t\varphi_{\tau}(z)$. Indeed, by Proposition 2.4, we have that

$$F_{\sigma_t}^{-1}(z) - z = t \cdot \varphi_{\sigma}(z) = t \cdot \varphi_{\tau}(F_{\rho}(z)),$$

and we thus obtain that

$$\begin{aligned} \varphi_{\mu_t}(F_{\rho}(z)) &= F_{\mu_t}^{-1}(F_{\rho}(z)) - F_{\rho}(z) \\ &= F_{\sigma_t}^{-1}(z) - F_{\rho}(z) \\ &= F_{\sigma_t}^{-1}(z) - z + z - F_{\rho}(z) \\ &= t \cdot \varphi_{\tau}(F_{\rho}(z)) + F_{\rho}^{-1}(F_{\rho}(z)) - F_{\rho}(z). \end{aligned}$$

By analytic continuation, we conclude that

$$\begin{aligned}\varphi_{\mu t}(z) &= t \cdot \varphi_\tau(z) + F_\rho^{-1}(z) - z \\ &= \varphi_\rho(z) + t \cdot \varphi_\tau(z),\end{aligned}$$

which completes the proof. $\square$

Remark 1. There are examples $\rho, \tau \in \mathcal{M}_\mathbb{R}$ such that $\tau \boxplus \rho \in \mathcal{ID}(\boxplus, \mathbb{R})$ but $\tau \notin \mathcal{ID}(\boxplus, \mathbb{R})$ and τ is not a summand of ρ , see [4].

Combining Proposition 2.4 and Lemma 2.5, we can reconstruct free convolution semigroups [27, 7] as follows.

Theorem 2.6. *Given $\mu \in \mathcal{M}_\mathbb{R}$, then $\mu^{\boxplus t} \in \mathcal{M}_\mathbb{R}$ is defined such that $\varphi_{\mu^{\boxplus t}}(z) = t\varphi_\mu(z)$ for all $t > 1$. Moreover, there exists an analytic map $\omega_t : \mathbb{C}^+ \rightarrow \mathbb{C}^+$ such that $F_{\mu^{\boxplus t}}(z) = F_\mu(\omega_t(z))$, for $z \in \mathbb{C}^+$, $\omega_t = F_{(\mu \boxplus \mu)^{\boxplus(t-1)}}(z)$ and $\varphi_{(\mu \boxplus \mu)^{\boxplus(t-1)}} = (t-1)(z - F_\mu(z))$ for all $t > 1$.*

Let $H_t(z) = z + (t-1)(z - F_\mu(z))$, then by Proposition 2.2 and Theorem 2.6, we know that H_t is the left inverse of ω_t such that $H_t(\omega_t(z)) = z$ for $z \in \mathbb{C}^+$. Therefore, for $t > 1$, $\omega_t(z)$ can be written as

$$(2.3) \quad \omega_t(z) = z + \left(1 - \frac{1}{t}\right) (F_{\mu^{\boxplus t}}(z) - z), \quad z \in \mathbb{C}^+.$$

We deduce from (2.3) and the definition of ω_t in Theorem 2.6 that, for $t > 0$,

$$z - F_{(\mu \boxplus \mu)^{\boxplus t}}(z) = \left(1 - \frac{1}{t+1}\right) (z - F_{\mu^{\boxplus(t+1)}}(z)),$$

which implies that

$$(2.4) \quad (\mu \boxplus \mu)^{\boxplus t}(z) = (\mu^{\boxplus(t+1)})^{\boxplus(t/(t+1))}.$$

2.3. Two formulas related to free Brownian motion. Given $\mu \in \mathcal{M}_\mathbb{R}$, we construct subordination functions ω_t as in Theorem 2.6. Let $\sigma_t = (\mu \boxplus \mu)^{\boxplus t} \in \mathcal{M}_\mathbb{R}$, then $\omega_{t+1} = F_{\sigma_t}(z)$ for $t > 0$. Given $\nu \in \mathcal{M}_\mathbb{R}$, let $\rho_t = \gamma_t \boxplus \nu$ and let $F_t = F_{\rho_t}(z)$ for all $t > 0$. From Proposition 2.4 and Theorem 2.6, we know that ρ_t and σ_t are $\boxplus$ -infinitely divisible and their Voiculescu transforms are given by $\varphi_{\rho_t}(z) = tG_\nu$ and $\varphi_{\sigma_t}(z) = t(z - F_\mu(z))$. By comparing Voiculescu transform of ρ_t with Voiculescu transform of σ_t , we deduce that $F_t = \omega_{t+1}$ for some $t > 0$ if and only if $G_\nu(z) = z - F_\mu(z)$.

For any $t > 0$, Belinschi and Nica [11] construct the transformation $\mathbb{B}_t : \mathcal{M}_\mathbb{R} \rightarrow \mathcal{M}_\mathbb{R}$ such that

$$\mathbb{B}_t(\mu) = (\mu^{\boxplus(1+t)})^{1/(1+t)}, \text{ for } \mu \in \mathcal{M}_\mathbb{R}.$$

They also show that $\mathbb{B}_t$ is a semigroup and $\mathbb{B}_2 = \mathbb{B}$, where the map $\mathbb{B} : \mathcal{ID}(\boxplus, \mathbb{R}) \rightarrow \mathcal{ID}(\boxplus, \mathbb{R})$ is the bijective map from the $\boxplus$ -infinitely divisible distributions to the $\boxplus$ -infinitely divisible distributions, discovered in the seminar paper [14]. The following theorem is from [11].

Theorem 2.7. *Let μ and ν be a pair of probability measures on the real line such that*

$$(2.5) \quad G_\nu(z) = z - F_\mu(z), \quad z \in \mathbb{C}^+.$$

Then we have

$$G_{\nu \boxplus \gamma_t}(z) = z - F_{\mathbb{B}_t(\mu)}(z), \quad t > 0, \quad z \in \mathbb{C}^+.$$

Remark 2. Given $\mu, \nu \in \mathcal{M}_{\mathbb{R}}$ satisfying (2.5), then Maassen [25] shows that μ has mean zero and variance one. Conversely, if $\mu \in \mathcal{M}_{\mathbb{R}}$ has mean zero and variance one, then there exists a unique $\nu \in \mathcal{M}_{\mathbb{R}}$ satisfying (2.5).

Given $\tau \in \mathcal{ID}(\boxplus, \mathbb{R})$ and $\mu, \nu \in \mathcal{M}_{\mathbb{R}}$, we compare free Lévy process $\nu \boxplus \tau^{\boxplus t}$ and free convolution semigroup $\mu^{\boxplus(t+1)}$. If $\varphi_\tau(F_\nu(z)) = z - F_\mu(z)$, then $\tau \boxplus \nu = \mu \boxplus \mu$, which implies that subordination function of $F_{\nu \boxplus (\tau^{\boxplus t})}$ to F_ν is the same as the subordination function of $F_{\mu^{\boxplus(t+1)}}$ to F_μ . The following theorem generalizes Theorem 2.7. The argument is similar to the proof of Theorem 1.6 in [11] (see also the proof of Lemma 3 in [4]), therefore we omit the proof.

Theorem 2.8. *Given $\tau \in \mathcal{ID}(\boxplus, \mathbb{R})$, and let μ and ν be a pair of probability measures on the real line such that*

$$\varphi_\tau(F_\nu(z)) = z - F_\mu(z), \quad z \in \mathbb{C}^+.$$

Then we have

$$\varphi_\tau(F_{\nu \boxplus (\tau^{\boxplus t})}(z)) = z - F_{\mathbb{B}_t(\mu)}(z), \quad t > 0, \quad z \in \mathbb{C}^+.$$

Remark 3. Let $\tau = \gamma_{a,b}$ be the semi-circular distribution with mean a and variance b , and let μ, ν be a pair of probability measures on the real line such that

$$(2.6) \quad \varphi_\tau(F_\nu(z)) = z - F_\mu(z).$$

We first compute

$$(2.7) \quad \begin{aligned} F_\mu(z) &= z - \varphi_\tau(F_\nu(z)) \\ &= z - \left(a + \frac{b}{z}\right) \circ F_\nu(z) \\ &= z - a - bG_\nu(z). \end{aligned}$$

By Theorem 2.8, then we have that

$$(2.8) \quad \begin{aligned} F_{\mathbb{B}_t(\mu)}(z) &= z - \varphi_\tau(F_{\nu \boxplus \tau^{\boxplus t}}(z)) \\ &= z - \left(\left(a + \frac{b}{z}\right) \circ F_{\nu \boxplus \gamma_{a,b}^{\boxplus t}}\right)(z) \\ &= z - a - bG_{\nu \boxplus \gamma_{a,b}^{\boxplus t}}(z). \end{aligned}$$

By (2.8) and the definition of Boolean convolution, we obtain that

$$(2.9) \quad F_{(\mathbb{B}_t(\mu))^{\boxplus t}}(z) = z - ta - tbG_{\nu \boxplus \gamma_{a,b}^{\boxplus t}}(z).$$

Equations (2.7), (2.8) and (2.9) were studied in [4]. We would like to point out that, as it was shown in [4, Proposition 1 and Example 1], $(\mathbb{B}_t(\mu))^{\boxplus t} \in \mathcal{ID}(\boxplus, \mathbb{R})$, and $(\mathbb{B}_t(\mu))^{\boxplus t} = (\tau \boxplus \nu)^{\boxplus t} = (\mu \boxplus \mu)^{\boxplus t}$. In fact, for all $\mu \in \mathcal{M}_{\mathbb{R}}$, we can deduce from (2.4) and the identity $(\mathbb{B}_t(\mu))^{\boxplus t} = (\mu^{\boxplus(1+t)})^{\boxplus t/(1+t)}$ that $(\mathbb{B}_t(\mu))^{\boxplus t}$ is the measure associated with the subordination function of $\mu^{\boxplus(1+t)}$ with respect to μ , that is $(\mathbb{B}_t(\mu))^{\boxplus t} = (\mu \boxplus \mu)^{\boxplus t}$.

3. MULTIPLICATIVE FREE CONVOLUTION AND MULTIPLICATIVE BOOLEAN CONVOLUTION ON $\mathcal{M}_{\mathbb{T}}$

Given any two probability measures μ, ν on $\mathbb{T}$, the unit circle of $\mathbb{C}$, we can define their multiplicative free convolution. We first recall the calculation of the multiplicative free convolution of two measures on $\mathbb{T}$ with nonzero means. Given $\mu \in \mathcal{M}_{\mathbb{T}}$, we define

$$\psi_{\mu}(z) = \int_{\mathbb{T}} \frac{tz}{1-tz} d\mu(t)$$

and set $\eta_{\mu}(z) = \psi_{\mu}(z)/(1 + \psi_{\mu}(z))$. The following proposition [8] characterizes the η -transforms of probability measures on $\mathbb{T}$.

Proposition 3.1. *Let $\eta : \mathbb{D} \rightarrow \mathbb{C}$ be an analytic function. Then the following assertions are equivalent.*

- (1) *There exists a probability measure $\mu \in \mathcal{M}_{\mathbb{T}}$ such that $\eta = \eta_{\mu}$.*
- (2) *$\eta(0) = 0$, and $|\eta(z)| < 1$ holds for all $z \in \mathbb{D}$.*

If $\mu \in \mathcal{M}_{\mathbb{T}} \cap \mathcal{M}_{*}$, then $\eta'_{\mu}(0) = \int_{\mathbb{T}} t d\mu(t) \neq 0$. Therefore, the inverse η_{μ}^{-1} is defined in a neighborhood of zero. We set $\Sigma_{\mu}(z) = \eta_{\mu}^{-1}(z)/z$. Given $\mu, \nu \in \mathcal{M}_{\mathbb{T}} \cap \mathcal{M}_{*}$, their multiplicative free convolution, which is denoted by $\mu \boxtimes \nu$, is the unique probability measure in $\mathcal{M}_{\mathbb{T}} \cap \mathcal{M}_{*}$ such that

$$(3.1) \quad \Sigma_{\mu \boxtimes \nu}(z) = \Sigma_{\mu}(z) \Sigma_{\nu}(z)$$

holds for z in a neighborhood of zero.

It is known [19, 9] that there exist two analytic functions $\omega_1, \omega_2 : \mathbb{D} \rightarrow \mathbb{D}$ such that

- (1) $\omega_1(0) = \omega_2(0) = 0$,
- (2) $\eta_{\mu \boxtimes \nu}(z) = \eta_{\mu}(\omega_1(z)) = \eta_{\nu}(\omega_2(z))$.

A probability measure $\mu \in \mathcal{M}_{\mathbb{T}}$ is said to be $\boxtimes$ -infinitely divisible if for any positive integer n , there exists $\mu_n \in \mathcal{M}_{\mathbb{T}}$ such that $\mu = \mu_n^{\boxtimes n} = \mu_n \boxtimes \cdots \boxtimes \mu_n$. It is shown in [15] that if $\mu \in \mathcal{M}_{\mathbb{T}} \setminus \mathcal{M}_{*}$ is $\boxtimes$ -infinitely divisible, then μ is the Haar measure on $\mathbb{T}$; and $\mu \in \mathcal{M}_{\mathbb{T}} \cap \mathcal{M}_{*}$ is $\boxtimes$ -infinitely divisible if and only if there exists a function

$$(3.2) \quad u(z) = \alpha i + \int_{\mathbb{T}} \frac{1+tz}{1-tz} d\sigma(t),$$

such that $\Sigma_{\mu}(z) = \exp(u(z))$, where $\alpha \in \mathbb{R}$ and σ is a finite positive measure on $\mathbb{T}$. Equation (3.2) is the analogue of the Lévy-Hinčin formula for multiplicative free

convolution on $\mathbb{T}$. The analogue of the normal distribution in this context is given by $\Sigma_{\lambda_t}(z) = \exp\left(\frac{t}{2} \frac{1+z}{1-z}\right)$. Denote by $\mathcal{ID}(\boxtimes, \mathbb{T})$ the set of all $\boxtimes$ -infinitely divisible measures on $\mathbb{T}$.

Lemma 3.2. *If $\mu \in \mathcal{M}_{\mathbb{T}} \cap \mathcal{M}_*$ is $\boxtimes$ -infinitely divisible. Then*

- (1) *The function $H(z) = z\Sigma_{\mu}(z)$ is the left inverse of $\eta_{\mu}(z)$, that is $H(\eta_{\mu}(z)) = z$ for all $z \in \mathbb{D}$.*
- (2) *The function η_{μ} extends to be a continuous function on $\overline{\mathbb{D}}$, and η_{μ} is one-to-one on $\overline{\mathbb{D}}$.*
- (3) *The set $\{z \in \mathbb{D} : |z\Sigma_{\mu}(z)| < 1\}$ is a simply connected domain which coincides with $\{\eta_{\mu}(z) : z \in \mathbb{D}\}$, and its boundary is $\eta_{\mu}(T)$ which is a simple closed curve.*

Proof. Observing that $H(\eta_{\mu}(z)) = z$ is valid in a neighborhood of zero, we obtain assertion (1) by analytic continuation.

Note that $H : \mathbb{D} \rightarrow \mathbb{C}$ satisfies the conditions in Proposition 4.5 in [8] and thus assertions (2) and (3) hold. $\square$

3.1. Multiplicative free Brownian motion. For $\mu \in \mathcal{M}_{\mathbb{T}}$ and $t > 0$, we study the multiplicative free convolution $\mu \boxtimes \lambda_t$. We first concentrate on the case when μ has nonzero mean. The case when μ has mean zero will be studied in Subsection 3.2.

We start with the following result which is the multiplicative version of Lemma 1 in [18].

Lemma 3.3. *Given $\mu, \nu \in \mathcal{M}_{\mathbb{T}} \cap \mathcal{M}_*$, we have that*

$$\eta_{\mu}(z) = \eta_{\mu \boxtimes \nu}(z \cdot \Sigma_{\nu}(\eta_{\mu}(z)))$$

holds for z in a neighborhood of zero.

Proof. From (3.1), we find that

$$(3.3) \quad \frac{\eta_{\mu \boxtimes \nu}^{-1}(z)}{z} = \frac{\eta_{\mu}^{-1}(z)}{z} \cdot \frac{\eta_{\nu}^{-1}(z)}{z}$$

holds for z in a neighborhood of zero, which we denote by D_0 . We choose a subdomain $D_1 \subset D_0$ such that $\eta_{\mu}(D_1) \subset D_0$. Replacing z by $\eta_{\mu}(z)$ in (3.3), we obtain that

$$(3.4) \quad \frac{\eta_{\mu \boxtimes \nu}^{-1}(\eta_{\mu}(z))}{\eta_{\mu}(z)} = \frac{\eta_{\mu}^{-1}(\eta_{\mu}(z))}{\eta_{\mu}(z)} \cdot \frac{\eta_{\nu}^{-1}(\eta_{\mu}(z))}{\eta_{\mu}(z)} = \frac{z}{\eta_{\mu}(z)} \cdot \frac{\eta_{\nu}^{-1}(\eta_{\mu}(z))}{\eta_{\mu}(z)}$$

holds for $z \in D_1$. Note that $\eta_{\nu}^{-1}(z) = z\Sigma_{\nu}(z)$ holds for $z \in D_0$, and we then rewrite (3.4) as

$$(3.5) \quad \eta_{\mu \boxtimes \nu}^{-1}(\eta_{\mu}(z)) = z\Sigma_{\nu}(\eta_{\mu}(z)).$$

Applying $\eta_{\mu \boxtimes \nu}$ on both sides of (3.5) yields that

$$\eta_{\mu}(z) = \eta_{\mu \boxtimes \nu}(z\Sigma_{\nu}(\eta_{\mu}(z)))$$

holds for z in a neighborhood of zero D_1 . $\square$

For any $t > 0$, we denote by $\eta_t : \mathbb{D} \rightarrow \mathbb{D}$ the subordination function of $\mu \boxtimes \lambda_t$ with respect to μ . Since $\eta_t : \mathbb{D} \rightarrow \mathbb{D}$ is analytic and $\eta_t(0) = 0$, Proposition 3.1 implies the existence of a probability measure ρ_t such that $\eta_{\rho_t}(z) = \eta_t(z)$.

Lemma 3.4. *The measure ρ_t is $\boxtimes$ -infinitely divisible and its Σ -transform is $\Sigma_{\rho_t}(z) = \Sigma_{\lambda_t}(\eta_{\mu}(z))$.*

Proof. Define analytic function $\Phi_t : \mathbb{D} \rightarrow \mathbb{C}$ by $\Phi_t(z) := z \Sigma_{\lambda_t}(\eta_{\mu}(z))$ for all $t > 0$. By Lemma 3.3, we have that

$$\eta_{\mu}(z) = \eta_{\mu \boxtimes \lambda_t}(z \Sigma_{\lambda_t}(\eta_{\mu}(z))) = \eta_{\mu \boxtimes \lambda_t}(\Phi_t(z))$$

which implies that

$$\eta_{\mu \boxtimes \lambda_t}(z) = \eta_{\mu}(\eta_t(z)) = \eta_{\mu \boxtimes \lambda_t}(\Phi_t(\eta_t(z))).$$

Since $\eta_{\mu \boxtimes \lambda_t}$ is invertible in a neighborhood of zero, we have that $\Phi_t(\eta_t(z)) = z$ in a neighborhood of zero.

We thus obtain that $\eta_{\rho_t}^{-1}(z) = \eta_t^{-1}(z) = \Phi_t(z)$ holds for z in a neighborhood of zero, which yields that

$$(3.6) \quad \Sigma_{\rho_t}(z) = \frac{\eta_{\rho_t}^{-1}(z)}{z} = \Sigma_{\lambda_t}(\eta_{\mu}(z)).$$

By the definition of the ψ - and η -transforms, we have that

$$(3.7) \quad \Sigma_{\lambda_t}(\eta_{\mu}(z)) = \exp \left(\frac{t}{2} \int_{\mathbb{T}} \frac{1 + \xi z}{1 - \xi z} d\mu(\xi) \right).$$

The real part of the integrand in (3.7) is positive for all $z \in \mathbb{D}$, then the asseration follows from (3.6) and Theorem 6.7 in [15]. $\square$

By (3.6), the right hand side of (3.7) is the Lévy-Hinčin representation of ρ_t . We can also write η_t in terms of λ_t and $\mu \boxtimes \lambda_t$. Replacing z by $\eta_{\mu \boxtimes \lambda_t}(z)$ in the equation

$$\frac{\eta_{\mu \boxtimes \lambda_t}^{-1}(z)}{z} = \frac{\eta_{\mu}^{-1}(z)}{z} \cdot \frac{\eta_{\lambda_t}^{-1}(z)}{z},$$

we obtain that

$$\frac{z}{\eta_{\mu \boxtimes \lambda_t}(z)} = \frac{\eta_t(z)}{\eta_{\mu \boxtimes \lambda_t}(z)} \cdot \Sigma_{\lambda_t}(\eta_{\mu \boxtimes \lambda_t}(z)),$$

which shows that

$$(3.8) \quad \eta_t(z) = \frac{z}{\Sigma_{\lambda_t}(\eta_{\mu \boxtimes \lambda_t}(z))}.$$

3.2. Modified $\mathcal{S}$ -transform and subordination functions. Given $\mu \in \mathcal{M}_T \setminus \mathcal{M}_*$ and $\nu \in \mathcal{M}_T \cap \mathcal{M}_*$, it is known from [19] that $\eta_{\mu \boxtimes \nu}$ is subordinated to η_μ and η_ν . The subordination function for this case is generally not unique. However, we show that there is a nice subordination function, which we call the principal subordination function, uniquely determined by certain conditions. Using the principal subordination function, results related to subordination function in the case $\mu, \nu \in \mathcal{M}_T \cap \mathcal{M}_*$ can be extended to the case where $\mu \in \mathcal{M}_T \setminus \mathcal{M}_*$ and $\nu \in \mathcal{M}_T \cap \mathcal{M}_*$.

Let us first give an example which illustrates the non-uniqueness of subordination functions.

Example 3.5. For $k \in \mathbb{N}$, and let $\lambda^{(k)} = 1/k \sum_{n=0}^{k-1} \delta_{z_n}$, where $z_n = e^{2\pi i n/k}$. We have $\psi_{\lambda^{(k)}}(z) = z^k/(1 - z^k)$ and $\eta_{\lambda^{(k)}}(z) = z^k$. Given $\nu \in \mathcal{M}_T \cap \mathcal{M}_*$, if $\omega : \mathbb{D} \rightarrow \mathbb{D}$ is a subordination function of $\eta_{\lambda^{(k)} \boxtimes \nu}$ with respect to $\eta_{\lambda^{(k)}}$, then $\omega^{(n)}(z) := e^{2\pi i n/k} \omega(z)$ is also a subordination function of $\eta_{\lambda^{(k)} \boxtimes \nu}$ with respect to $\eta_{\lambda^{(k)}}$ for all integer $0 < n < k$.

We now introduce the modified $\mathcal{S}$ -transform. Given two free random variables x and y in a W^* -probability space $(\mathcal{A}, \phi)$, such that $\phi(x) = 0$ and $\phi(y) \neq 0$. We can not directly apply Voiculescu's $\mathcal{S}$ -transform (Σ -transform) to calculate the distribution of xy . N. Raj Rao and R. Speicher [28] introduce a new transform, which we call the modified $\mathcal{S}$ -transform, to deal with this case. They apply the modified $\mathcal{S}$ -transform to study the distribution of xy where x, y are free self-adjoint random variables such that $\phi(x) = 0$, $\phi(y) \neq 0$. For nonzero self-adjoint operator x , we have that $\phi(x^2) \neq 0$.

Assume that $\phi(x) = \dots = \phi(x^{k-1}) = 0$ and $\phi(x^k) \neq 0$, Arizmendi [5] observe that we can calculate the distribution of xy using the idea in [28]. We present the details of their work for reader's convenience.

We first recall some definitions. For $\mu \in \mathcal{M}_T \cap \mathcal{M}_*$, we have $\psi_\mu(0) = 0$ and $\psi'_\mu(0) \neq 0$. It follows that there exists a function $\chi_\mu(z)$, which is analytic in a neighborhood of zero, such that

$$\psi_\mu(\chi_\mu(z)) = \chi_\mu(\psi_\mu(z)) = z$$

holds for sufficiently small z . The usual $\mathcal{S}$ -transform is defined by

$$\mathcal{S}_\mu(z) = \frac{z+1}{z} \chi_\mu(z).$$

We then have

$$\Sigma_\mu(z) = \mathcal{S}_\mu\left(\frac{z}{1-z}\right), \quad \eta_\mu^{-1}(z) = \chi\left(\frac{z}{1-z}\right).$$

We set

$$\mathcal{M}_T^k = \{\mu \in \mathcal{M}_T : \int_T t^n d\mu(t) = 0 \text{ for } 1 \leq n < k, \text{ and } \int_T t^k d\mu(t) \neq 0\}.$$

Then for $\mu \in \mathcal{M}_T^k$, we have that

$$(3.9) \quad \begin{cases} \psi'_\mu(0) = \cdots \psi_\mu^{(k-1)}(0) = 0 = \eta'_\mu(0) = \cdots \eta_\mu^{(k-1)}(0), \\ \psi_\mu^{(k)}(0) \neq 0 \text{ and } \eta_\mu^{(k)} \neq 0. \end{cases}$$

For $\mu \in \mathcal{M}_T^k, \nu \in \mathcal{M}_T \cap \mathcal{M}_*$, from the definition of free independence, we deduce that $\mu \boxtimes \nu \in \mathcal{M}_T^k$.

We recall the following classical result in complex analysis (see, for example, [23]).

Theorem 3.6. *If $f(z)$ is holomorphic in $|z| < R$, and suppose that*

$$f(0) = f'(0) = \cdots = f^{(k-1)}(0) = 0, \quad f^{(k)} \neq 0,$$

then for small values of $w \neq 0$ the equation

$$f(z) = w$$

has k roots $z_1(w), \dots, z_k(w)$, which tend to zero when w tends to zero. Moreover, there exists a function $g(w)$, holomorphic for w sufficiently small with $g(0) = 0$ and $g'(0) \neq 0$, such that for any fixed small values $w \neq 0$,

$$z_j(w) = g(\omega^j w^{1/k}), \quad \omega = e^{2\pi i/k}, \quad 0 \leq \arg \omega^{1/k} < \frac{2\pi}{k},$$

if we put those roots in a certain order.

Remark 4. The converse of Theorem 3.6 is also true. More precisely, if we are given a function $g(w)$ which is holomorphic for w sufficiently small with $g(0) = 0$ and $g'(0) \neq 0$, and for $j = 1, \dots, k$, let

$$z_j(w) = g(\omega^j w^{1/k}), \quad \omega = e^{2\pi i/k}, \quad 0 \leq \arg \omega^{1/k} < \frac{2\pi}{k},$$

then $z_1(w), \dots, z_k(w)$ are the roots of the equation

$$F^k(z) = w,$$

where F is a holomorphic function defined in a neighborhood of the zero such that $F(g(w)) = w$.

For $j = 1, \dots, k$, denote $D_{j,r} = \{\omega^j z : 0 \leq \arg(z) < 2\pi/k, |z| < r\}$. We record the following result for convenience.

Proposition 3.7. *Under the assumption of Theorem 3.6, we have that $z_j(f(z)) = z$ for $z \in g(D_{j,r})$ for r sufficiently small.*

Given $\mu \in \mathcal{M}_T^k$, and by Theorem 3.6, we know that there exist k functions represented by the power series in $z^{1/k}$ such that

$$(3.10) \quad \psi_\mu(\chi_\mu^{(j)}(z)) = z,$$

for z sufficiently small. Moreover, there exists a function $g_\mu(w)$ holomorphic in a neighborhood of the zero, such that for $j = 1, \dots, k$,

$$\chi_\mu^{(j)}(z) = g_\mu(\omega^j z^{1/k}),$$

where $\omega = e^{2\pi i/k}$, $0 \leq \arg z^{1/k} < 2\pi/k$.

Definition 3.8. Given $\mu \in \mathcal{M}_T^k$. Let $\chi_\mu^{(j)}$ be the inverse function of ψ_μ in (3.10), the modified $\mathcal{S}$ -transform of μ is k functions $\mathcal{S}_\mu^{(1)}(z), \dots, \mathcal{S}_\mu^{(k)}(z)$, such that for $j = 1, \dots, k$,

$$\mathcal{S}_\mu^{(j)}(z) = \chi_\mu^{(j)}(z) \cdot \frac{1+z}{z}.$$

Given $\mu \in \mathcal{M}_T^k$ and $\nu \in \mathcal{M}_T \cap \mathcal{M}_*$, we set

$$\mathcal{S}^{(j)}(z) = \mathcal{S}_\mu^{(j)}(z) \cdot \mathcal{S}_\nu(z)$$

and compute

$$\begin{aligned} \chi^{(j)}(z) &= \mathcal{S}^{(j)}(z) \cdot \frac{z}{1+z} \\ &= \mathcal{S}_\mu^{(j)}(z) \cdot \mathcal{S}_\nu(z) \cdot \frac{z}{1+z} \\ (3.11) \quad &= \chi_\mu^{(j)}(z) \cdot \mathcal{S}_\nu(z) \\ &= g_\mu(\omega^j z^{1/k}) \cdot \mathcal{S}_\nu(z) \\ &= g(\omega^j z^{1/k}), \end{aligned}$$

where $g(z) = g_\mu(z) \cdot \mathcal{S}_\nu(z^k)$ is a function such that $g(0) = 0$, $g'(0) \neq 0$. From Remark 4, we deduce that for different j , there exists the same left inverse ψ such that $\psi(\chi^{(j)}(z)) = z$. Therefore, we have the following proposition.

Proposition 3.9. Given $\mu \in \mathcal{M}_T^k$ and $\nu \in \mathcal{M}_T \cap \mathcal{M}_*$, for $1 \leq j \leq k$, let

$$\begin{aligned} \mathcal{S}^{(j)}(z) &= \mathcal{S}_\mu^{(j)}(z) \cdot \mathcal{S}_\nu(z) \\ \chi^{(j)}(z) &= \mathcal{S}^{(j)}(z) \cdot \frac{z}{1+z}. \end{aligned}$$

Then there exists a unique holomorphic function ψ defined in a neighborhood of the zero such that

$$\psi((\chi^{(j)})(z)) = z.$$

The following result is due to Raj Rao and Speicher [28] and Arizmendi [5].

Theorem 3.10. Given $\mu \in \mathcal{M}_T^k, \nu \in \mathcal{M}_T \cap \mathcal{M}_*$, we have that

$$(3.12) \quad \mathcal{S}_{\mu \boxtimes \nu}^{(j)}(z) = \mathcal{S}_\mu^{(j)}(z) \cdot \mathcal{S}_\nu(z), \quad j = 1, \dots, k,$$

where the modified $\mathcal{S}$ -transforms are listed in a certain order.

Because of Proposition 3.9 and Theorem 3.10, for fixed $\mu \in \mathcal{M}_T^k$ and $\nu \in \mathcal{M}_T \cap \mathcal{M}_*$, we denote

$$(3.13) \quad \psi(z) = \psi_{\mu \boxtimes \nu}(z), \text{ and } \chi^{(j)}(z) = \chi_{\mu \boxtimes \nu}^{(j)}(z),$$

and we also denote $g(z) = g_\mu(z) \cdot \mathcal{S}_\nu(z^k)$ as in (3.11).

Given $\mu \in \mathcal{M}_T^k, \nu \in \mathcal{M}_T \cap \mathcal{M}_*$, we set $\iota_\mu^{(j)}(z) = \chi_\mu^{(j)}(z/(1-z))$ and $\iota_\nu(z) = \chi_\nu(z/(1-z))$. Theorem 3.10 implies that

$$(3.14) \quad \chi_{\mu \boxtimes \nu}^{(j)}(z) = \chi_\mu^{(j)}(z) \cdot \chi_\nu(z) \cdot \frac{1+z}{z}.$$

We also have that $\chi_\mu^{(j)}(z) = g_\mu(\omega^j z^{1/k})$ and $\chi_{\mu \boxtimes \nu}^{(j)}(z) = g_\mu(\omega^j z^{1/k}) \cdot \mathcal{S}_\nu(z) = g(\omega^j z^{1/k})$. Substituting z by $\psi_{\mu \boxtimes \nu}(z)$ in (3.14), and applying Proposition 3.7, we find that

$$z = \chi_{\mu \boxtimes \nu}^{(j)}(\psi_{\mu \boxtimes \nu}(z)) = \chi_\mu^{(j)}(\psi_{\mu \boxtimes \nu}(z)) \cdot \chi_\nu(\psi_{\mu \boxtimes \nu}(z)) \cdot \frac{1 + \psi_{\mu \boxtimes \nu}(z)}{\psi_{\mu \boxtimes \nu}(z)},$$

where $z \in g(D_{j,r})$ for r sufficiently small. We thus have that

$$(3.15) \quad z\eta_{\mu \boxtimes \nu} = \iota_\mu^{(j)}(\eta_{\mu \boxtimes \nu}) \cdot \iota_\nu(\eta_{\mu \boxtimes \nu})$$

holds in the same domain.

We can now utilize the argument in [9] to prove the existence of subordination function of $\eta_{\mu \boxtimes \nu}$ with respect to η_μ for $\mu \in \mathcal{M}_T^k, \nu \in \mathcal{M}_T \cap \mathcal{M}_*$. Note that part of the following result is known in [19].

Theorem 3.11. *Given $\mu \in \mathcal{M}_T^k, \nu \in \mathcal{M}_T \cap \mathcal{M}_*$, there exists two unique analytic functions $\omega_1, \omega_2 : \mathbb{D} \rightarrow \mathbb{D}$ such that*

- (1) $\omega_1(0) = \omega_2(0) = 0$;
- (2) $\eta_{\mu \boxtimes \nu}(z) = \eta_\mu(\omega_1(z)) = \eta_\nu(\omega_2(z))$,
- (3) $\omega_1(z)\omega_2(z) = z\eta_{\mu \boxtimes \nu}(z)$ for all $z \in \mathbb{D}$.

Proof. Since $\eta_\mu(0) = 0, \eta_\nu(0) = 0$, we can write $\eta_\mu(z) = zf_1(z), \eta_\nu(z) = zf_2(z)$ for two analytic functions $f_1, f_2 : \mathbb{D} \rightarrow \mathbb{D}$. Fix $1 \leq j \leq k$, set $\omega_1(z) = \iota_\mu^{(j)}(\eta_{\mu \boxtimes \nu}(z))$, $\omega_2(z) = \iota_\nu(\eta_{\mu \boxtimes \nu}(z))$ defined in $g(D_{j,r})$ for r sufficiently small.

By (3.15), we have that

$$z\eta_{\mu \boxtimes \nu}(z) = \omega_1(z)\omega_2(z),$$

holds for $z \in g(D_{j,r})$. We thus obtain that

$$\omega_1(z) = \frac{z\eta_{\mu \boxtimes \nu}}{\omega_2(z)} = \frac{z\eta_\nu(\omega_2(z))}{\omega_2(z)} = zf_2(\omega_2(z)).$$

Similarly, we have that $\omega_2(z) = zf_1(\omega_1(z))$ for $z \in g(D_{j,r})$. Regarding $\omega_1(z), \omega_2(z)$ as Denjoy-Wolff points, the same argument in [9] implies ω_1, ω_2 can be extended analytically to $\mathbb{D}$. By the uniqueness of Denjoy-Wolff points, ω_1, ω_2 does not depend on the choice of j .

By the definitions of $\iota_\mu^{(j)}, \iota_\nu$, we have that $\eta_\mu(\omega_1(z)) = \eta_\nu(\omega_2(z)) = \eta_{\mu \boxtimes \nu}$ for z in $g(D_{j,r})$. Thus (2) and (3) hold by analytic continuation. Since $\eta'_\nu(0) \neq 0$, η_ν is locally invertible near the origin and therefore ω_2 is unique. Finally (3) implies the uniqueness of ω_1 . $\square$

Since $\mu \in \mathcal{M}_T^k$ and $\mu \boxtimes \nu \in \mathcal{M}_T^k$, we have that $\omega'_1(0) \neq 0$, where ω_1 is given in Theorem 3.11.

Definition 3.12. For $\mu \in \mathcal{M}_T^k, \nu \in \mathcal{M}_T \cap \mathcal{M}_*$, the subordination ω_1 satisfying the relations (1), (2) and (3) in Theorem 3.11 is called the *principal* subordination function of $\eta_{\mu \boxtimes \nu}$ with respect to η_μ . The measure $\rho \in \mathcal{M}_T \cap \mathcal{M}_*$ satisfying $\eta_\rho(z) = \omega_1(z)$ is called the *principal* subordination distribution of $\eta_{\mu \boxtimes \nu}$ with respect to η_μ .

Note that for $\mu, \nu \in \mathcal{M}_T \cap \mathcal{M}_*$, the principal subordination function of $\eta_{\mu \boxtimes \nu}$ with respect to η_μ is the usual subordination function.

The following result might be obtained by approximation. We provide a direct proof.

Corollary 3.13. *Given $\mu \in \mathcal{M}_T^k, \nu \in \mathcal{M}_T \cap \mathcal{M}_*$, let ρ be the principal subordination distribution of $\eta_{\mu \boxtimes \nu}$ with respect to η_μ . Then we have that*

$$\Sigma_\rho(z) = \Sigma_\nu(\eta_\mu(z)).$$

In particular, if $\nu \in \mathcal{ID}(\boxtimes, T)$, then $\rho \in \mathcal{ID}(\boxtimes, T)$.

Proof. By choosing a sequence $\mu_n \in \mathcal{M}_T \cap \mathcal{M}_*$ such that μ_n converges to μ weakly, Lemma 3.3 implies

$$\eta_\mu(z) = \eta_{\mu \boxtimes \nu}(z \Sigma_\nu(\eta_\mu(z)))$$

for z in a neighborhood of zero.

Set $\Phi(z) = z \Sigma_\nu(\eta_\mu(z)) = z \cdot \mathcal{S}_\nu(\psi_\mu(z))$, and we thus have

$$(3.16) \quad \eta_{\mu \boxtimes \nu}(z) = \eta_\mu(\omega_1(z)) = \eta_{\mu \boxtimes \nu}(\Phi(\omega_1(z))) = \eta_{\mu \boxtimes \nu}(\Phi(\eta_\rho(z))).$$

Fix $1 \leq j \leq k$, we claim that if $z \in g(D_{j,r})$, then $\Phi(\omega_1(z)) = z$. Indeed, for $0 \leq \arg(w^{1/k}) < 2\pi/k$ and $z = g(\omega^j w^{1/k}) = \chi^{(j)}(w)$, using the construction of ω_1 in Theorem 3.11, we have

$$(3.17) \quad \omega_1(z) = \iota_\mu^{(j)}(\eta_{\mu \boxtimes \nu}(z)) = \chi_\mu^{(j)}(\psi_{\mu \boxtimes \nu}(z)).$$

From (3.11) and (3.13), we have

$$(3.18) \quad \chi^{(j)}(w) = g(\omega^j w^{1/k}), \text{ and } \psi_{\mu \boxtimes \nu}(\chi^{(j)}(w)) = w.$$

Equations (3.17) and (3.18) imply that $\omega_1(g(\omega^j w^{1/k})) = \chi_\mu^{(j)}(w)$.

Note that $\psi_\mu(\omega_1(g(\omega^j w^{1/k}))) = \psi_{\mu \boxtimes \nu}(g(\omega^j w^{1/k})) = w$. Thus we obtain that

$$\Phi(\omega_1(z)) = \Phi(\omega_1(g(\omega^j w^{1/k}))) = \chi_\mu^{(j)}(w) \mathcal{S}_\nu(w) = \chi^{(j)}(w) = z.$$

The above claim, (3.16) and Proposition 3.7 imply

$$z = \Phi(\omega_1(z)) = \Phi(\eta_\rho(z))$$

for $z \in g(D_{j,r})$. We conclude that $\Sigma_\rho(z) = \Phi(z)/z = \Sigma_\nu(\eta_\mu(z))$ for z in a small neighborhood of zero by applying the above argument for all $1 \leq j \leq k$.

If $\nu \in \mathcal{ID}(\boxtimes, \mathbb{T})$, then by Theorem 6.7 in [16], there exists an analytic function $u(z)$ defined in $\mathbb{D}$ such that $\Sigma_\nu(z) = \exp(u(z))$ and $\Re u(z) \geq 0$ for all $z \in \mathbb{D}$. Thus $\Sigma_\nu(\eta_\mu(z)) = \exp(u(\eta_\mu(z)))$ and $\Re(u(\eta_\mu(z))) \geq 0$ for all $z \in \mathbb{D}$, and then the second assertion follows from Theorem 6.7 in [16]. $\square$

Remark 5. If $k = 1$, noticing that $\mathcal{M}_T^k = \mathcal{M}_T \cap \mathcal{M}_*$, the modified $\mathcal{S}$ -transform is the usual $\mathcal{S}$ -transform. We see that Corollary 3.14 holds when $\mu, \nu \in \mathcal{M}_T \cap \mathcal{M}_*$.

The following result is the multiplicative analogue of Lemma 2.6.

Proposition 3.14. *Given $\rho, \tau \in \mathcal{M}_T \cap \mathcal{M}_*$, let σ be a measure in $\mathcal{M}_T \cap \mathcal{M}_*$ such that $\eta_{\rho \boxtimes \tau}(z) = \eta_\rho(\eta_\sigma(z))$. If $\sigma \in \mathcal{ID}(\boxtimes, \mathbb{T})$, then $\rho \boxtimes \tau^{\boxtimes t}$ can be defined for all $t \geq 0$ in the sense that $\Sigma_{\rho \boxtimes (\tau^{\boxtimes t})}(z) = \Sigma_\rho(z)(\Sigma_\tau(z))^t$.*

Proof. For $t > 0$, there exists $\mu_t \in \mathcal{M}_T \cap \mathcal{M}_*$ such that

$$\eta_{\mu_t}(z) = \eta_\rho(\eta_{\sigma^{\boxtimes t}}(z)).$$

Using a similar argument as in the proof of Lemma 2.6 and applying Corollary 3.13, we can find that $\Sigma_{\mu_t}(z) = \Sigma_\rho(z)(\Sigma_\tau(z))^t$. $\square$

3.3. Semigroups related to multiplicative free convolution. Recall that $\mathcal{M}_T^* = \{\mu \in \mathcal{M}_T \cap \mathcal{M}_* : \eta_\mu(z) \neq 0, \forall z \in \mathbb{D} \setminus \{0\}\}$. Given $\mu \in \mathcal{M}_T^*$ and $t > 1$, and let u be an analytic function satisfying that $z/(\eta_\mu(z)) = e^{u(z)}$ holds for z in a neighborhood of zero. Set $H_t(z) = ze^{(t-1)u(z)} = z[z/(\eta_\mu(z))]^{t-1}$. It is shown in [8] that H_t has a right inverse $\omega_t : \mathbb{D} \rightarrow \mathbb{D}$ such that $H_t(\omega_t(z)) = z$, and there exists a probability measure $\mu^{\boxtimes t} \in \mathcal{M}_T^*$ such that

- (1) $\eta_{\mu^{\boxtimes t}}(z) = \eta_\mu(\omega_t(z))$ and $\Sigma_{\mu^{\boxtimes t}}(z) = (\Sigma_\mu(z))^t$,
- (2) $\omega_t(z) = \eta_{\mu^{\boxtimes t}}(z) [z/\eta_{\mu^{\boxtimes t}}(z)]^{1/t}$ for $z \in \mathbb{D}$, where the power is chosen such that the equation holds.

Observe that for each $t > 0$, by Proposition 3.1, there exists a probability measure $\sigma_t \in \mathcal{M}_T$ such that $\eta_{\sigma_t}(z) = \omega_{t+1}(z)$. It turns out that σ_t is $\boxtimes$ -infinitely divisible and its Σ -transform is $\Sigma_{\sigma_t}(z) = [z/\eta_\mu(z)]^t$, which can be obtained by applying the same argument as in the proof of Lemma 3.4.

The following result is a partial converse of Theorem 3.5 in [8].

Theorem 3.15. *Given $\mu \in \mathcal{M}_T \cap \mathcal{M}_*$, assume that for any $t > 1$, there exists a probability measure $\mu_t \in \mathcal{M}_T$ such that*

$$(3.19) \quad \Sigma_{\mu_t}(z) = (\Sigma_\mu(z))^t.$$

Assume in addition that μ_t is subordinated with respect to μ for all $t > 1$. Then $\eta_\mu(z) \neq 0$ for all $z \in \mathbb{D} \setminus \{0\}$, that is $\mu \in \mathcal{M}_T^*$.

Proof. For each $t > 1$, we denote by ω_t the subordination function of μ_t to μ . Observing that $\mu_t \in \mathcal{M}_*$ and $\omega'_t(0) \neq 0$, for each $t > 1$, there exists a probability measure $\sigma_{t-1} \in \mathcal{M}_T \cap \mathcal{M}_*$ such that $\eta_{\sigma_{t-1}}(z) = \omega_t(z)$. We rewrite (3.19) as

$$(3.20) \quad \frac{\eta_{\mu_t}^{-1}(z)}{z} = \left[\frac{\eta_{\mu}^{-1}(z)}{z} \right]^t$$

for z in a neighborhood of zero.

Note that $\omega_t^{-1}(z) = \eta_{\mu_t}^{-1}(\eta_{\mu}(z))$ for z in a neighborhood of zero. Replacing z by $\eta_{\mu}(z)$ in (3.20), we obtain that

$$\begin{aligned} \frac{\omega_t^{-1}(z)}{\eta_{\mu}(z)} &= \frac{\eta_{\mu_t}^{-1}(\eta_{\mu}(z))}{\eta_{\mu}(z)} \\ &= \left[\frac{\eta_{\mu}^{-1}(\eta_{\mu}(z))}{\eta_{\mu}(z)} \right]^t \\ &= \left[\frac{z}{\eta_{\mu}(z)} \right]^t, \end{aligned}$$

which implies that,

$$\frac{\omega_t^{-1}(z)}{z} = \left[\frac{z}{\eta_{\mu}(z)} \right]^{t-1}.$$

Given $t > 0$, we thus have $\Sigma_{\sigma_t}(z) = [z/\eta_{\mu}(z)]^t$ for z in a neighborhood of zero. Therefore σ_t is $\boxtimes$ -infinitely divisible.

By Theorem 6.7 in [15], there exists an analytic function $u(z)$ in $\mathbb{D}$ such that $\Re u(z) \geq 0$ if $z \in \mathbb{D}$ and $\Sigma_{\sigma_1}(z) = \exp(u(z))$. We thus obtain that $z/\eta_{\mu}(z) = \exp(u(z))$, which implies that $\eta_{\mu}(z) \neq 0$ for all $z \in \mathbb{D}$. $\square$

It was pointed in [8] that $\mu^{\boxtimes t}$ is only determined up to a rotation by a multiple of $2\pi t$. Note that ω_t and σ_t are determined by the choice of $\mu^{\boxtimes t}$.

3.4. Multiplicative Boolean convolution and the Bercovici-Pata bijection.

Multiplicative Boolean convolution on T was studied by Franz [22]. Let $\mu \in \mathcal{M}_T$, and we set $k_{\mu}(z) = z/\eta_{\mu}(z)$. Given two probability measures $\mu, \nu \in \mathcal{M}_T$, their multiplicative Boolean convolution $\mu \boxtimes \nu$ is a probability measure on T such that

$$k_{\mu \boxtimes \nu}(z) = k_{\mu}(z)k_{\nu}(z)$$

for all $z \in \mathbb{D}$.

A probability $\mu \in \mathcal{M}_T$ is said to be $\boxtimes$ -infinitely divisible, if for any positive integer n , there exists $\mu_n \in \mathcal{M}_T$ such that $\mu = \mu_n^{\boxtimes n}$. Let P_0 be the Haar measure. It is

shown in [22] that $\mu \in \mathcal{M}_T \setminus \{P_0\}$ is $\boxtimes$ -infinitely divisible if and only if $\eta'_\mu(0) \neq 0$ and $\eta_\mu \neq 0$ for all $z \in \mathbb{D} \setminus \{0\}$, that is $\mu \in \mathcal{M}_T^*$, which is equivalent to

$$(3.21) \quad k_\mu(z) = \exp \left(bi + \int_T \frac{1 + \xi z}{1 - \xi z} d\tau_\mu(\xi) \right),$$

where $b \in \mathbb{R}$ and τ_μ is a finite measure on T . Equation (3.21) is the analogue of the Lévy-Hinčin formula in this context.

The multiplicative Bercovici-Pata bijection from $\boxtimes$ to $\boxtimes$ was studied in [31]. Denote the set of all $\boxtimes$ -infinitely divisible measures on T by $\mathcal{ID}(\boxtimes, T)$, and the multiplicative Bercovici-Pata bijection from $\boxtimes$ to $\boxtimes$ by $\mathbb{M}$. Then we have $k_\mu(z) = \Sigma_{\mathbb{M}(\mu)}(z)$.

Given $\mu \in \mathcal{ID}(\boxtimes, T) \setminus P_0 = \mathcal{M}_T^*$, let ω_2 be the subordination function of $\mu^{\boxtimes 2}$ with respect to μ , and let σ be the probability measure on T such that $\eta_\sigma(z) = \omega_2(z)$. Then σ is $\boxtimes$ -infinitely divisible and its Σ -transform is $\Sigma_\sigma(z) = z/\eta_\mu(z) = k_\mu(z)$. Therefore, σ is the same as $\mathbb{M}(\mu)$. Since $P_0 \boxtimes P_0 = P_0$ and $\eta_{P_0} = z$, the subordination function of $P_0 \boxtimes P_0$ with respect to P_0 is the identity map z , and the measure associated with the identity map z is P_0 . To summarize, we have the following corollary.

Corollary 3.16. *Given $\mu \in \mathcal{ID}(\boxtimes, T)$, let ω_2 be the subordination function of $\mu^{\boxtimes 2}$ with respect to μ , and let σ be the probability measure on T such that $\eta_\sigma(z) = \omega_2(z)$. Then $\sigma = \mathbb{M}(\mu)$, where $\mathbb{M}$ is the multiplicative Bercovici-Pata bijection from $\boxtimes$ to $\boxtimes$.*

Proposition 3.17. *Let $\mu \in \mathcal{M}_T$, then the following are equivalent.*

- (1) $\mu \in \mathcal{ID}(T, \boxtimes)$;
- (2) $\delta_\beta \boxtimes \mu \in \mathcal{ID}(T, \boxtimes)$ for any $\beta \in T$.

Proof. It is enough to prove that (1) implies (2) for $\mu \in \mathcal{M}_T \cap \mathcal{M}_*$. Observing that $\eta_{\delta_\beta \boxtimes \mu}(z) = \beta \cdot \eta_\mu(z)$, we thus have

$$\Sigma_{\delta_\beta \boxtimes \mu}(z) = \frac{\eta_{\delta_\beta \boxtimes \mu}^{-1}(z)}{z} = \frac{\eta_\mu^{-1}(\bar{\beta}z)}{z} = \bar{\beta} \cdot \Sigma_\mu(\bar{\beta}z).$$

The result follows from the Lévy-Hinčin formula for the multiplicative free convolution on T . $\square$

3.5. An analogue of equations studied by Belinschi and Nica. In this subsection, we prove our Theorem 1.1. Recall that λ_t is the free multiplicative analogue of the normal distribution on T , the unit circle of $\mathbb{C}$, with $\Sigma_{\lambda_t}(z) = \exp\left(\frac{t}{2} \frac{1+z}{1-z}\right)$ and we set $\lambda = \lambda_1$. For $\mu \in \mathcal{M}_T$, we denote $m_1(\mu) = \int_T \xi d\mu(\xi)$.

Proposition 3.18. *Given $\mu \in \mathcal{M}_T^*$, and an analytic map $u = u(z)$ defined by*

$$(3.22) \quad u(z) = bi + \int_T \frac{1 + \xi z}{1 - \xi z} d\tau(\xi), \quad z \in \mathbb{D},$$

with $b \in [0, 2\pi)$ and τ a finite measure on T . If $k_\mu(z) = z/\eta_\mu(z) = \exp(u(z))$, then $b = \arg(1/m_1(\mu)) \in [0, 2\pi)$, and $\tau(T) = \ln |1/m_1(\mu)|$. In particular, there exists a probability measure $\nu \in \mathcal{M}_T$ such that $k_\mu(z) = \Sigma_\lambda(\eta_\nu(z))$ if and only if $m_1(\mu) = e^{-1/2}$.

Proof. By definition, we have that

$$(3.23) \quad k_\mu(0) = \lim_{z \rightarrow 0} \frac{z}{\eta_\mu(z)} = \frac{1}{\eta'_\mu(0)} = \frac{1}{m_1(\mu)}.$$

Since $u(0) = bi + \tau(T)$, we then obtain that

$$(3.24) \quad b = \arg\left(\frac{1}{m_1(\mu)}\right), \text{ and } \tau(T) = \ln\left|\frac{1}{m_1(\mu)}\right|.$$

The first assertion follows.

By (3.7), we have that

$$\Sigma_\lambda(\eta_\nu(z)) = \exp\left(\frac{1}{2} \int_T \frac{1 + \xi z}{1 - \xi z} d\nu(\xi)\right).$$

Noticing that k_μ has the Herglotz representation as (3.21), we conclude that $k_\mu(z)$ can be written in the form of $\Sigma_\lambda(\eta_\nu(z))$ for a probability measure ν on T if and only if $\ln(1/m_1(\mu)) = 1/2$. This implies the second half of the assertion. $\square$

For $\mu \in \mathcal{ID}(\boxtimes, T) \setminus P_0 = \mathcal{M}_T^*$ with $m_1(\mu) > 0$, let $u(z)$ be the analytic function satisfying $k_\mu(z) = \exp(u(z))$ and $u(0) > 0$. Given $t > 1$, let $H_t(z) = z \exp((t-1)u(z))$, and denote its right inverse by $\omega_t : \mathbb{D} \rightarrow \mathbb{D}$ with $\omega_t(0) = 0$. We define (see [8]) $\mu^{\boxtimes t}$ by the relation

$$(3.25) \quad \eta_{\mu^{\boxtimes t}}(z) = \eta_\mu(\omega_t(z)).$$

Then we see that $H'_t(0) > 0$, $\omega'_t(0) > 0$ and that

$$m_1(\mu^{\boxtimes t}) = \eta'_{\mu^{\boxtimes t}}(0) > 0.$$

For $t > 0$, we also define $\mu^{\boxtimes t}$ by the relation

$$(3.26) \quad k_{\mu^{\boxtimes t}}(z) = \exp(tu(z)).$$

For this choice of the Boolean convolution power, we have that

$$m_1(\mu^{\boxtimes t}) > 0.$$

Definition 3.19. Given $\mu \in \mathcal{M}_T^*$ such that $m_1(\mu) > 0$, we define a family of maps $\{\mathbb{M}_t\}_{t \neq 0}$ by

$$\mathbb{M}_t(\mu) = \left(\mu^{\boxtimes(t+1)} \right)^{\boxtimes \frac{1}{t+1}},$$

where we choose $\mu^{\boxtimes(t+1)}$ and $\mathbb{M}_t(\mu)$ in a way such that they have positive means.

The next result is a special case of Theorem 4.4 in [6].

Lemma 3.20. *Given $\mu \in \mathcal{M}_T^*$ with $m_1(\mu) > 0$, then the following assertions are true.*

- (1) $\mathbb{M}_{t+s}(\mu) = \mathbb{M}_t(\mathbb{M}_s(\mu))$ for all $t, s \geq 0$.
- (2) $\mathbb{M}_1(\mu) = \mathbb{M}(\mu)$.

We are now able to prove Theorem 1.1.

Proof of Theorem 1.1. We set

$$(3.27) \quad u(z) = \frac{1}{2} \int_T \frac{1 + \xi z}{1 - \xi z} d\nu(\xi),$$

then by (3.7) and the assumption (1.2), we have that

$$(3.28) \quad \frac{z}{\eta_\mu(z)} = \Sigma_\lambda(\eta_\nu(z)) = \exp(u(z)).$$

By Proposition 3.18, we see that $m_1(\mu) > 0$. We therefore can choose the multiplicative convolution power $\mu^{\boxtimes(t+1)}$ such that $m_1(\mu^{\boxtimes(t+1)}) > 0$.

Let η_t be the principal subordination function of $\nu \boxtimes \lambda_t$ with respect to ν and ω_{t+1} be the subordination function of $\mu^{\boxtimes(t+1)}$ with respect to μ . Let $\rho_t, \sigma_t \in \mathcal{M}_T$ such that $\eta_{\rho_t} = \eta_t$ and $\eta_{\sigma_t} = \omega_{t+1}$.

By Corollary 3.13, (1.2) implies that $\Sigma_{\rho_t}(z) = \Sigma_{\lambda_t}(\eta_\mu(z)) = \exp(tu(z))$. From the choice of $\mu^{\boxtimes(t+1)}$, the function $H_{t+1}(z) := z \exp(tu(z))$ is the left inverse of ω_{t+1} such that $H_{t+1}(\omega_{t+1}(z)) = z$ for all $z \in \mathbb{D}$, which implies that

$$(3.29) \quad \Sigma_{\sigma_t}(z) = \exp(tu(z)).$$

We thus obtain that $\rho_t = \sigma_t$ and $\eta_t = \omega_{t+1}$.

Replacing z by η_t in (1.2), we obtain that

$$(3.30) \quad \begin{aligned} \Sigma_\lambda(\eta_{\nu \boxtimes \lambda_t}(z)) &= \Sigma_\lambda(\eta_\nu((\eta_t(z)))) = \frac{\eta_t(z)}{\eta_\mu(\eta_t(z))} \\ &= \frac{\omega_{t+1}(z)}{\eta_\mu(\omega_{t+1}(z))} = \frac{\omega_{t+1}(z)}{\eta_{\mu^{\boxtimes(t+1)}}(z)} \\ &= \left(\frac{z}{\eta_{\mu^{\boxtimes(t+1)}}(z)} \right)^{1/t+1}. \end{aligned}$$

On the other hand, by the definition of $\mathbb{M}_t$, we have that

$$\begin{aligned} \frac{z}{\eta_{\mathbb{M}_t(\mu)}(z)} &= \frac{z}{\eta_{(\mu^{\boxtimes(t+1)})^{\boxtimes \frac{1}{t+1}}}(z)} \\ &= \left(\frac{z}{\eta_{\mu^{\boxtimes(t+1)}}(z)} \right)^{1/t+1}. \end{aligned}$$

We thus complete the proof of Theorem 1.1. $\square$

3.6. Some examples and applications. We start with some examples which are the multiplicative analogue of examples studied in [1], [4], [6] and [11]. We define the set

$$(A) = \{\mu \in \mathcal{M}_T^* : m_1(\mu) = e^{-1/2}\}.$$

By Proposition 3.22, the set $\mathcal{M}_T$ is in one-to-one correspondence with the set (A) via the bijection $\nu \leftrightarrow \mu$, such that $\Sigma_\lambda(\eta_\nu(z)) = z/\eta_\mu(z)$.

Definition 3.21. The bijective map $\Lambda : \mathcal{M}_T \rightarrow (A)$ is defined by

$$\Sigma_\lambda(\eta_\nu(z)) = \frac{z}{\eta_{\Lambda[\nu]}(z)}, \quad \forall \nu \in \mathcal{M}_T.$$

Using the notation Λ , Theorem 1.1 implies that

$$\Lambda[\nu \boxtimes \lambda_t] = \mathbb{M}_t[\Lambda(\nu)], \quad \forall \nu \in \mathcal{M}_T.$$

Example 3.22. Let δ_1 be the Dirac measure at 1, and let $\mu = \Lambda[\delta_1]$, then we have $z/\eta_\mu(z) = \Sigma_\lambda(\eta_{\delta_1}(z)) = \exp(\frac{1}{2} \frac{1+z}{1-z})$. For $t \geq 0$, Theorem 1.1 implies that

$$\frac{z}{\eta_{\mathbb{M}_t(\mu)}(z)} = \Sigma_\lambda(\eta_{\delta_1 \boxtimes \lambda_t}(z)) = \Sigma_\lambda(\eta_{\lambda_t}(z)).$$

In particular, when $t = 1$,

$$\eta_{\mathbb{M}_1(\mu)}(z) = \frac{z}{\Sigma_\lambda(\eta_{\lambda_1}(z))} = \eta_{\lambda_1}(z),$$

where we used the equality $((z\Sigma_\lambda) \circ \eta_\lambda)(z) = z$ and $\lambda = \lambda_1$. Therefore, $\mathbb{M}_1(\mu)$ is the free multiplicative analogue of the normal distribution on T .

Example 3.23. More generally, we consider $\lambda_{b,t} = \delta_b \boxtimes \lambda_t$ and $\mu_{b,t} = \Lambda[\lambda_{b,t}]$. Then we have that

$$\Sigma_\lambda(\eta_{\lambda_{b,t}}(z)) = \frac{z}{\eta_{\mu_{b,t}}(z)}, \quad \text{for } t \neq 0.$$

On the other hand, Theorem 1.1 implies that

$$\Sigma_\lambda(\eta_{\lambda_{b,t_1+t_2}}(z)) = \Sigma_\lambda(\eta_{\lambda_{b,t_1} \boxtimes \lambda_{t_2}}(z)) = \frac{z}{\eta_{\mathbb{M}_{t_2}(\mu_{b,t_1})}(z)}, \quad \text{for } t_1, t_2 \geq 0,$$

which yields that $\mu_{b,t_1+t_2} = \mathbb{M}_{t_2}(\mu_{b,t_1})$ for $t_1, t_2 \geq 0$.

We would like to provide another example which covers part of Example 4.10 in [6].

Example 3.24. Let P_0 be the Haar measure on $\mathbb{T}$. Then by the free independence, $P_0 \boxtimes \lambda_t = P_0$. We set $\mu = \Lambda[P_0]$, and then we have

$$\Sigma_\lambda(\eta_{P_0 \boxtimes \lambda_t}(z)) = \Sigma_\lambda(\eta_{P_0}(z)) = \frac{z}{\eta_\mu(z)},$$

which implies that $\mathbb{M}_t(\mu) = \mu$ for all $t \geq 0$. To calculate the distribution of μ , we note that $\eta_{P_0} \equiv 0$, which shows that $\eta_\mu = e^{-1}z$, and we thus have that $\psi_\mu(z) = z/e - z$. Using the identity

$$\frac{1}{\pi} \left(\psi_\mu(z) + \frac{1}{2} \right) = \frac{1}{2\pi} \int_0^{2\pi} \frac{e^{it} + z}{e^{it} - z} d\mu(e^{-it}),$$

and Stieltjes's inversion formula, we obtain that

$$\mu(dt) = \frac{1}{2\pi} \frac{1 - e^{-2}}{1 + e^{-2} - 2e^{-1} \cos(t)} dt, \quad 0 \leq t \leq 2\pi.$$

We then give some applications of results concerning infinity divisibility of the measures associated with subordination functions. For $\mu \in \mathcal{M}_\mathbb{T}$, we say μ is nontrivial if it is not a Dirac measure at a point on $\mathbb{T}$.

Lemma 3.25. *Given $\sigma \in \mathcal{ID}(\boxtimes, \mathbb{T})$ which is non-trivial, and $0 < \epsilon < 1$, there exists a positive number $n(\epsilon)$ such that*

$$\eta_{\sigma_t}(\overline{\mathbb{D}}) \subset \mathbb{D}_\epsilon = \{z = re^{i\theta} : 0 \leq r \leq \epsilon, 0 \leq \theta < 2\pi\}$$

holds for any $t > n(\epsilon)$, where $\sigma_t = \sigma^{\boxtimes t}$.

Proof. If $\sigma = P_0$, the Haar measure on $\mathbb{T}$, then the result is trivial. If $\sigma \neq P_0$ is nontrivial, then by Theorem 6.7 in [15], there exists a finite positive measure ν on $\mathbb{T}$ with $\nu(\mathbb{T}) > 0$, $\alpha \in \mathbb{R}$, and an analytic function u defined by

$$u(z) = i\alpha + \int_{\mathbb{T}} \frac{1 + \xi z}{1 - \xi z} d\nu(\xi), \text{ for } z \in \mathbb{D},$$

such that $\Sigma_\sigma(z) = \exp(u(z))$. We choose $\sigma_t \in \mathcal{M}_\mathbb{T}$ satisfying $\Sigma_{\sigma_t}(z) = \exp(tu(z))$. Noticing that other choices of the multiplicative free convolution power of σ can be obtained from σ_t by a rotation of a multiple of $2\pi t$, it is enough to prove the assertion for σ_t .

We set $\Phi_{\sigma_t} = z\Sigma_{\sigma_t}(z)$, then by Lemma 3.2, we have that

$$\Phi_{\sigma_t}^{-1}(\mathbb{D}) = \eta_{\sigma_t}(\mathbb{D}).$$

For $z = re^{i\theta} \in \mathbb{D}$, we calculate

$$\begin{aligned}
 |\Phi_{\sigma_t}(z)| &= r \exp \left(t \int_{\mathbb{T}} \frac{1-r^2}{|1-\xi z|^2} d\nu(\xi) \right) \\
 &\geq r \exp \left(t \int_{\mathbb{T}} \frac{1-r^2}{|1+r|^2} d\nu(\xi) \right) \\
 &= r \exp \left(t \cdot \nu(\mathbb{T}) \frac{1-r}{1+r} \right).
 \end{aligned}
 \tag{3.31}$$

Since $\lim_{t \rightarrow \infty} r \exp \left(t \cdot \nu(\mathbb{T}) \frac{1-r}{1+r} \right) = \infty$, we deduce that for any $0 < \epsilon < 1$, there exists a positive number $n(\epsilon)$ such that, for all $t > n(\epsilon)$, we have that

$$|\Phi_{\sigma_t}(z)| > 1, \text{ for } |z| = \epsilon.$$

By Lemma 3.2, $\Phi_{\sigma_t}(\mathbb{D})$ is a simply connected domain which contains zero, which implies that

$$\eta_{\sigma_t}(\mathbb{D}) = \Phi_{\sigma_t}^{-1}(\mathbb{D}) \subset \mathbb{D}_\epsilon, \text{ for } t > n(\epsilon).$$

The assertion follows by the fact that η_{σ_t} extends to be a continuous function on $\overline{\mathbb{D}}$. $\square$

For $\mu \in \mathcal{M}_{\mathbb{T}}$, we have that

$$\frac{1}{2\pi} \left(\frac{1 + \eta_\mu(z)}{1 - \eta_\mu(z)} \right) = \frac{1}{2\pi} \int_0^{2\pi} \frac{e^{i\theta} + z}{e^{i\theta} - z} d\mu(e^{-i\theta}), \quad z \in \mathbb{D}.$$

The real part of this function is the Poisson integral of the measure $d\mu(e^{-i\theta})$, we can recover μ by Stieltjes's inversion formula. The functions

$$\frac{1}{2\pi} \Re \left(\frac{1 + \eta_\mu(re^{i\theta})}{1 - \eta_\mu(re^{i\theta})} \right) = \frac{1}{2\pi} \frac{1 - |\eta_\mu(re^{i\theta})|^2}{|1 - \eta_\mu(re^{i\theta})|^2}$$

converge to the density of $\mu(e^{-i\theta})$ a.e. relative to Lebesgue measure, and they converge to infinity a.e. relative to the singular part of this measure.

Proposition 3.26. *Given $\mu \in \mathcal{M}_{\mathbb{T}}$ and $\sigma \in \mathcal{ID}(\boxtimes, \mathbb{T})$ which is nontrivial, let μ_t be the unique probability measure on $\mathbb{T}$ such that*

$$\eta_{\mu_t}(z) = \eta_\mu(\eta_{\sigma_t}(z)).$$

Then we have

$$\lim_{t \rightarrow \infty} \sup_{\theta \in [0, 2\pi]} \left| \frac{d\mu_t(e^{i\theta})}{d\theta} - \frac{1}{2\pi} \right| = 0,$$

where $d\mu_t(e^{i\theta})/d\theta$ is the density function of μ_t at $e^{i\theta}$ with respect to Lebesgue measure.

Proof. Given $0 < \epsilon < 1$, by Lemma 3.25, there exists $n(\epsilon) > 0$ such that $\eta_{\sigma_t}(e^{i\theta}) < \epsilon$ for $t \geq n(\epsilon)$, which yields that $\eta_{\mu_t}(z)$ extends continuously to $\overline{\mathbb{D}}$. We thus have that

$$|\eta_{\mu_t}(e^{i\theta})| = |\eta_\mu(\eta_{\sigma_t}(e^{i\theta}))| \leq |\eta_{\sigma_t}(e^{i\theta})| < \epsilon,$$

which implies that

$$(3.33) \quad \frac{1-\epsilon}{1+\epsilon} = \frac{1-\epsilon^2}{|1+\epsilon|^2} \leq \frac{1-|\eta_{\mu_t}(e^{i\theta})|^2}{|1-\eta_{\mu_t}(e^{i\theta})|^2} \leq \frac{1}{|1-\epsilon|^2}.$$

Since ϵ is arbitrary, combining (3.32) with (3.33), we prove our assertion. $\square$

Corollary 3.27. *Given $\mu \in \mathcal{M}_T$ and a nontrivial measure $\nu \in \mathcal{ID}(\boxtimes, T)$, then the density functions of the measures $\mu \boxtimes \nu_t$ converge to $1/2\pi$ uniformly as $t \rightarrow \infty$; if $\mu \in \mathcal{M}_T$ is nontrivial, then the density functions of the measures $\mu^{\boxtimes t}$ converge to $1/2\pi$ uniformly as $t \rightarrow \infty$.*

Proof. Noticing Corollary 3.13, Propositions 3.14–3.26 and Subsection 3.3, we only need to prove the case of $\mu^{\boxtimes t}$ for $\mu \notin \mathcal{M}_T^*$. We point out that the measures are nontrivial imply that the subordination distributions involved are nontrivial.

For $\mu \in \mathcal{M}_T \setminus \mathcal{M}_*$, we have $\mu^{\boxtimes n} = P_0$, where P_0 is the Haar measure on T . Thus the assertion is true for this case. For $\mu \in \mathcal{M}_T \cap \mathcal{M}_*$, but $\mu \notin \mathcal{M}_T^*$, it is shown in [8] that $\mu \boxtimes \mu \in \mathcal{M}_T^*$, thus this case reduces to the case when $\mu \in \mathcal{M}_T^*$. This finishes the proof. $\square$

4. MULTIPLICATIVE CONVOLUTION ON $\mathcal{M}_{\mathbb{R}^+}$

4.1. Multiplicative free convolution on $\mathcal{M}_{\mathbb{R}^+}$. We are interested in the probability measures on the positive real line $\mathbb{R}^+$, which are different from the Dirac measure at zero, we thus set

$$\mathcal{M}_{\mathbb{R}^+}^* = \mathcal{M}_{\mathbb{R}^+} \setminus \{\delta_0\}.$$

Given $\mu \in \mathcal{M}_{\mathbb{R}^+}^*$, we define

$$\psi_\mu(z) = \int_0^{+\infty} \frac{tz}{1-tz} d\mu(t),$$

and $\eta_\mu(z) = \psi_\mu(z)/(1 + \psi_\mu(z))$. The transform η_μ is characterized by the following proposition (see [8]).

Proposition 4.1. *Let $\eta : \mathbb{C} \setminus \mathbb{R}^+ \rightarrow \mathbb{C}$ be an analytic function such that $\eta(\bar{z}) = \overline{\eta(z)}$ for all $z \in \mathbb{C} \setminus \mathbb{R}^+$. Then the following two conditions are equivalent.*

- (1) $\eta = \eta_\mu$ for some $\mu \in \mathcal{M}_{\mathbb{R}^+}^*$.
- (2) $\eta(0-) = 0$ and $\arg(\eta(z)) \in [\arg z, \pi)$ for all $z \in \mathbb{C}^+$.

It can be shown that η_μ is invertible in some neighborhood of $(-\infty, 0)$, and we set $\Sigma_\mu(z) = \eta_\mu^{-1}(z)/z$ where η_μ^{-1} is defined in some neighborhood of $(\alpha, 0)$. Given two measures $\mu, \nu \in \mathcal{M}_{\mathbb{R}^+}^*$, the multiplicative free convolution of μ and ν is the probability measure $\mu \boxtimes \nu$ in $\mathcal{M}_{\mathbb{R}^+}^*$ such that

$$\Sigma_{\mu \boxtimes \nu}(z) = \Sigma_\mu(z) \Sigma_\nu(z)$$

holds in some neighborhood of $(\alpha, 0)$, where these functions are defined.

It is known from [19] and [9] that there exist two analytic functions $\omega_1, \omega_2 : \mathbb{C} \setminus \mathbb{R}^+ \rightarrow \mathbb{C} \setminus \mathbb{R}^+$ such that

- (1) $\omega_j(0-) = 0$ for $j = 1, 2$,
- (2) for any $\lambda \in \mathbb{C}^+$, we have $\omega_j(\bar{\lambda}) = \overline{\omega_j(\lambda)}$ for $j = 1, 2$,
- (3) $\eta_{\mu \boxtimes \nu}(z) = \eta_\mu(\omega_1(z)) = \eta_\nu(\omega_2(z))$ for $z \in \mathbb{C} \setminus \mathbb{R}^+$.

For simplicity, we say that ω_1 (resp. ω_2) is the subordination function of $\mu \boxtimes \nu$ with respect to μ (resp. ν), and $\mu \boxtimes \nu$ is subordinated to μ and ν .

The analogy of the Lévy-Hinčin in this setting was proved in [15] and [16]. Given $\mu \in \mathcal{M}_{\mathbb{R}^+}$, then $\mu \in \mathcal{M}_{\mathbb{R}^+}^*$ is $\boxtimes$ -infinitely divisible if and only if $\Sigma_\mu(z) = \exp(u(z))$, with

$$u(z) = a - bz + \int_0^{+\infty} \frac{1+tz}{z-t} d\sigma(t),$$

where $b \in \mathbb{R}$ and σ is a finite positive measure on $\mathbb{R}^+$. The analogue of the normal distribution in this context is given by $\Sigma_{\lambda_t}(z) = \exp\left(\frac{t}{2} \frac{z+1}{z-1}\right)$.

Lemma 4.2. *Let $\mu, \nu \in \mathcal{M}_{\mathbb{R}^+}^*$, then we have*

$$\eta_\mu(z) = \eta_{\mu \boxtimes \nu}(z \Sigma_\nu(\eta_\mu(z)))$$

holds in some neighborhood of interval $(\alpha, 0)$.

The proof of Lemma 4.2 is identical to the proof of Lemma 3.3, therefore we omit the details.

For any $t > 0$, assume that $\eta_t : \mathbb{D} \rightarrow \mathbb{D}$ is the subordination function of $\mu \boxtimes \lambda_t$ with respect to μ , by Lemma 4.2 and the characterization of η -transform, there exists a probability measure ρ_t in $\mathcal{M}_{\mathbb{R}^+}^*$ such that $\eta_{\rho_t}(z) = \eta_t(z)$. The argument in the proof of Lemma 3.4 implies the following result.

Proposition 4.3. *The measure ρ_t is $\boxtimes$ -infinitely divisible and its Σ -transform is $\Sigma_{\rho_t}(z) = \Sigma_{\lambda_t}(\eta_\mu(z))$, and*

$$\Sigma_{\lambda_t}(\eta_\mu(z)) = \exp\left(\frac{t}{2} \int_0^{+\infty} \frac{1+\xi z}{\xi z - 1} d\mu(\xi)\right).$$

We now discuss free convolution semigroups. Given $t > 1$, it is proved in [8] that one can define $\mu^{\boxtimes t} \in \mathcal{M}_{\mathbb{R}^+}^*$ such that $\Sigma_{\mu^{\boxtimes t}}(z) = (\Sigma_\mu(z))^t$ for $z < 0$ sufficiently close to zero. Similar to the case of $\mathcal{M}_{\mathbb{T}}$, $\mu^{\boxtimes t}$ is subordinated with respect to μ and we denote the subordination function by ω_t . By Theorem 2.6 in [8] and the characterization of η -transform, there exists a probability $\sigma_t \in \mathcal{M}_{\mathbb{R}^+}^*$ such that $\eta_{\sigma_t}(z) = \omega_{t+1}$ for all $t > 0$. Moreover, σ_t is $\boxtimes$ -infinitely divisible and its Σ -transform is $\Sigma_{\sigma_t}(z) = [z/\eta_\mu(z)]^t$.

4.2. Multiplicative Boolean convolution on $\mathcal{M}_{\mathbb{R}^+}$ and the semigroup $\mathbb{M}_t$. Bercovici proved in [13] that the multiplicative Boolean convolution does not preserve

$\mathcal{M}_{\mathbb{R}^+}$. But we can still define $\mu^{\boxtimes t}$ for $\mu \in \mathcal{M}_{\mathbb{R}^+}$ and $0 \leq t \leq 1$ as follows. Let $k_\mu(z) = z/\eta_\mu(z)$, the Boolean convolution power $\mu^{\boxtimes t}$ is defined by

$$k_{\mu^{\boxtimes t}}(z) = (k_\mu(z))^t.$$

The following definition was given in [6].

Definition 4.4. A family of maps from $\mathcal{M}_{\mathbb{R}^+}$ to itself is defined by

$$\mathbb{M}_t(\mu) = (\mu^{\boxtimes(t+1)})^{\boxtimes \frac{1}{t+1}}.$$

It is also shown in [6] that $\mathbb{M}_{t+s} = \mathbb{M}_t \circ \mathbb{M}_s$ for $t, s \geq 0$.

4.3. Analogous equations. Given a pair of probability measures $\nu, \mu \in \mathcal{M}_{\mathbb{R}^+}$, we also consider, as in the case $\mathcal{M}_{\mathbb{T}}$, the semigroups $\nu \boxtimes \lambda_t$ and $\mu^{\boxtimes(t+1)}$, the subordination functions η_t and ω_{t+1} , and their associated probability measures ρ_t, σ_t for all $t > 0$. Since $\Sigma_{\rho_t}(z) = \Sigma_{\lambda_t}(\eta_\nu(z))$ and $\Sigma_{\sigma_t}(z) = [z/\eta_\mu(z)]^t$, we deduce that $\eta_t = \omega_{t+1}$ if and only if

$$\Sigma_\lambda(\eta_\nu(z)) = \frac{z}{\eta_\mu(z)}.$$

Applying the same argument as in the proof of the Theorem 1.1, we obtain the following result.

Theorem 4.5. *Given a pair of probability measures $\mu, \nu \in \mathcal{M}_{\mathbb{R}^+}$, such that*

$$(4.1) \quad \Sigma_\lambda(\eta_\nu(z)) = \frac{z}{\eta_\mu(z)}, z \in \mathbb{C}^+.$$

Then we have

$$\Sigma_\lambda(\eta_{\nu \boxtimes \lambda_t}(z)) = \frac{z}{\eta_{\mathbb{M}_t(\mu)}(z)}, z \in \mathbb{C}^+.$$

5. A DESCRIPTION OF THE ANALOGUE OF THE NORMAL DISTRIBUTION

In this section, we describe the density functions of the free multiplicative analogue of the normal distributions.

5.1. The circle case. Let $\lambda_t \in \mathcal{M}_{\mathbb{T}}$ be the analogue of the normal distribution such that $\Sigma_{\lambda_t}(z) = \exp(\frac{t}{2} \frac{1+z}{1-z})$. We set $\Phi_t(z) = z\Sigma_{\lambda_t}(z)$, and let $\Omega_t = \{z \in \mathbb{D} : |\Phi_t(z)| < 1\}$. By Lemma 3.2, η_{λ_t} extends continuously to the unit circle $\mathbb{T}$, Ω_t is simply connected and bounded by a simple closed curve, and we have that $\partial\Omega_t = \eta_{\lambda_t}(\mathbb{T})$.

Observe that for $t \neq 4$, Φ_t has zeros of order one at $z_1(t) = (2 - t + \sqrt{t^2 - 4t})/2$ and $z_2(t) = (2 - t - \sqrt{t^2 - 4t})/2$. Φ_4 has a zero of order two at -1 ; and for all t , Φ_t has an essential singularity at 1 , and no other zeros and singularities. For $0 < t < 4$, $z_1(t), z_2(t) \in \mathbb{T}$ and $z_2(t) = \overline{z_1(t)}$, we let $\theta_1(t) \in (0, \pi)$ and $\theta_2(t) \in (\pi, 2\pi)$ such that $z_1(t) = e^{i\theta_1(t)}$ and $z_2(t) = e^{i\theta_2(t)}$. We have $z_1(4) = z_2(4) = -1$ and for $t > 4$, $z_1(t) \in (-1, 0)$ and $z_2(t) \in (-\infty, -1)$.

We define

$$g_t(r, \theta) = r \exp\left(\frac{t}{2} \frac{1 - r^2}{1 - 2r \cos \theta + r^2}\right) = |\Phi_t(z)|$$

for $z = re^{i\theta}$. The unit circle is parametrized by $T = \{e^{i\theta} : 0 \leq \theta < 2\pi\}$.

Lemma 5.1. *For $0 < t < 4$, $\partial\Omega_t = \{z = e^{i\theta} : \theta_1(t) \leq \theta \leq \theta_2(t)\} \cup \mathcal{L}_{1,t} \cup \mathcal{L}_{2,t}$, where $\mathcal{L}_{1,t}$ is an analytic curve, and $\mathcal{L}_{1,t}$ is in $\mathbb{D} \cap \mathbb{C}^+$ except one of its endpoints, and $\mathcal{L}_{2,t}$ is the reflection of $\mathcal{L}_{1,t}$ about x -axis. $\mathcal{L}_{1,t}$ can be parametrized by $\gamma_t(u)$ ($0 \leq u \leq 1$) such that $\gamma_t(0) \in \mathbb{R}$, $\gamma_t(1) = z_1(t)$ and $\gamma_t(u) \subset \mathbb{D} \cap \mathbb{C}^+$ for $0 < u < 1$. Moreover, $|\gamma_t(u)|$ is an increasing function of u on the interval $[0, 1]$.*

Proof. Observing that $\Phi_t(\bar{z}) = \overline{\Phi_t(z)}$, we see that $\partial\Omega_t$ is symmetric with respect to x -axis. Since Ω_t is simply connected and $\partial\Omega_t$ is a simple closed curve, $\partial\Omega_t$ intersects x -axis at two points.

Restricting Φ_t to real numbers, we find that $\Phi_t(\mathbb{R}) \subset \mathbb{R}$, and that Φ_t is an increasing function on $(-1, 1)$ since $\Phi'_t(z)$ is positive for $z \in (-1, 1)$. From $\Phi_t(-1) = -1$ and $\lim_{z \rightarrow 1^-} \Phi_t(z) = +\infty$, we deduce that

$$(5.1) \quad \Phi_t^{-1}((-1, 1)) = (-1, x(t)),$$

where $x(t)$ is the unique solution of the equation $\Phi_t(z) = 1$ for $z \in (-1, 1)$. The fact that $\Phi'_t(z) \neq 0$ for $z \neq z_1(t), z_2(t)$ implies that Φ_t is locally invertible for $z \neq z_1(t), z_2(t)$. Combining the fact that $\Phi_t(T \setminus \{1\}) \subset T$, we obtain that

$$\{e^{i\theta} : \theta_1(t) \leq \theta \leq \theta_2(t)\} \subset \partial\Omega_t$$

and $\partial\Omega_t$ has corners of opening $\pi/2$ at $z_1(t)$ and $z_2(t)$.

Since Φ_t is a conformal mapping from Ω_t to $\mathbb{D}$, by the symmetry $\Phi_t(\bar{z}) = \overline{\Phi_t(z)}$ and (5.1), noting that $\Phi'_t(0) = 1$, we thus deduce that $\Phi_t(\Omega_t \cap \mathbb{C}^+) \subset \overline{\mathbb{D}} \cap \mathbb{C}^+$. $\partial\Omega_t$ is a simple closed curve, thus $z_1(t)$ and $x(t)$ are connected by $\partial\Omega_t$. It is clear that $\partial\Omega_t \setminus \{e^{i\theta} : \theta_1(t) \leq \theta \leq \theta_2(t)\}$ does not intersect with T , we thus assume the curve $\gamma_t = \{\gamma_t(u) : 0 \leq u \leq 1\}$ is the part of Ω_t which connects $z_1(t)$ and $x(t)$ such that $\gamma_t(0) = x(t)$, $\gamma_t(1) = z_1(t)$ and $\gamma_t(u) \in \mathbb{D}$ for $0 < u < 1$.

We claim that $|\gamma_t(u)|$ is an increasing function of u on the interval $[0, 1]$. For given $0 < r < 1$, we define the function of θ by

$$g_{t,r}(\theta) = g_t(r, \theta) = |\Phi_t(re^{i\theta})|.$$

Then $g_{t,r}$ is a strictly decreasing function of θ on the interval $[0, \pi]$. From the fact that Ω_t is simply connected, we deduce that, for $z_0 \in \overline{\Omega_t} \cap \mathbb{D} \cap \mathbb{C}^+$, the arc

$$(5.2) \quad \{re^{i\theta} : |r| = |z_0|, \arg z_0 < \theta \leq \pi\} \subset \Omega_t.$$

Given $0 < u_1 < u_2 < 1$, we need to prove that $|\gamma_t(u_1)| < |\gamma_t(u_2)|$. Since $[0, x(t)] \subset \overline{\Omega_t}$, we obtain from (5.2) that

$$(5.3) \quad \{re^{i\theta} : 0 \leq r \leq x(t), 0 < \theta \leq \pi\} \subset \Omega_t,$$

which shows that $|\gamma_t(u_1)| > x(t)$. Suppose that $|\gamma_t(u_1)| \geq |\gamma_t(u_2)|$, then there exists $0 < u'_1 \leq u_1$ such that $|\gamma_t(u'_1)| = |\gamma_t(u_2)|$. If $\arg(\gamma_t(u'_1)) > \arg(\gamma_t(u_2))$, then by (5.2), $\gamma_t(u_2) \in \Omega_t$ and thus $\gamma_t(u_2) \notin \partial\Omega_t$; if $\arg(\gamma_t(u'_1)) < \arg(\gamma_t(u_2))$, then $\gamma_t(u'_1) \in \Omega_t$ and thus $\gamma_t(u'_1) \notin \partial\Omega_t$. For both cases, we obtain a contradiction. Thus $|\gamma_t(u_1)| < |\gamma_t(u_2)|$ and our claim is proved. $\square$

For $t > 0$, we let $x_1(t) \in (0, 1)$ be the unique solution of the equation $\Phi_t(z) = 1$ for $z \in (0, 1)$. For $0 < t \leq 4$ we let $x_2(t) = -1$; for $t > 4$, we let $x_2(t) \in (-1, 0)$ be the unique solution of the equation $\Phi_t(z) = -1$ for $z \in (-1, 0)$.

Lemma 5.2. *For $t \geq 4$, $\partial\Omega_t = \mathcal{L}_{1,t} \cup \mathcal{L}_{2,t}$, where $\mathcal{L}_{1,t}$ is an analytic curve, and $\mathcal{L}_{1,t}$ is in $\mathbb{D} \cap \mathbb{C}^+$ except its endpoints, and $\mathcal{L}_{2,t}$ is the reflection of $\mathcal{L}_{1,t}$ about x -axis. $\mathcal{L}_{1,t}$ can be parametrized by $\gamma_t(u)$ ($0 \leq u \leq 1$) such that $\gamma_t(0) = x_1(t)$, $\gamma_t(1) = x_2(t)$ and $\gamma_t(u) \subset \mathbb{D} \cap \mathbb{C}^+$ for $0 < u < 1$. Moreover, $|\gamma_t(u)|$ is an increasing function of u on the interval $[0, 1]$.*

Proof. Recall that Φ_4 has a zero of order two at -1 . For all $t > 4$, $z_2(t) < -1$ and $z_1 \in (-1, 0)$. The assertion follows from the similar arguments in the proof Lemma 5.1. $\square$

From the proof of Lemmas 5.1 and 5.2, for $t > 0$, we have that $\Phi_t^{-1}((-1, 1)) = (x_2(t), x_1(t))$. Moreover, $x_1(t) = \min\{|z| : z \in \partial\Omega_t\}$ and $-x_2(t) = \max\{|z| : z \in \partial\Omega_t\}$.

Remark 6. In fact, for any $t > 0$, from the equation

$$g_t(r, \theta) = 0, \quad 0 < r < 1, \quad 0 \leq \theta \leq \pi,$$

we can prove that $dr/d\theta > 0$ for $0 < \theta < \pi$, which implies that if $z \in \partial\Omega_t$, then the entire radius $\{rz : 0 \leq r < 1\}$ is contained in Ω_t . Therefore, $\partial\Omega_t$ can be parametrized by θ .

Lemma 5.3. *Using the same notations in Lemmas 5.1 and 5.2, for $t > 0$, the function $|1 - \gamma_t(u)|$ is an increasing function of u on $[0, 1]$.*

Proof. We only prove the case when $0 < t < 4$, the proof for other cases are similar. Noticing that $|1 - re^{i\theta}|^2 = 1 - 2\cos\theta + r^2$, since $|\gamma_t(u)|$ is an increasing function of u , to prove the assertion, we only need to prove that for the implicit function $r \exp(\frac{t}{2} \frac{1-r^2}{h}) = 1$ of r and h , then h increases when r increases on $(0, 1)$. From this equation, we have $h = h(r) = -(t/2)(1 - r^2)/(\ln r)$. One can check that $h'(r) > 0$ for $0 < r < 1$, therefore h is an increasing function of r . $\square$

Theorem 5.4. *Denote by A_t the support of λ_t .*

- (1) *For $t > 0$, the measure λ_t has no singular part, and its density function is an analytic function. $A_{t_1} \subset A_{t_2}$ if $t_1 < t_2 < 4$. $A_t \subsetneq \mathbb{T}$ for $0 < t < 4$ and $A_t = \mathbb{T}$ for $t \geq 4$.*
- (2) *The measure λ_t is unimodal for all $t > 0$ and its density is maximal at $z = 1$ and is minimal at $z = -1$.*

(3) The density function $d\lambda_t/d\theta$ converges uniformly to $1/(2\pi)$ as $t \rightarrow \infty$.

Proof. Since $z = 1$ is not in the closure of $\Omega_t = \eta_{\lambda_t}(\mathbb{D})$, then λ_t has no singular part. From the analyticity of Φ_t and or a general theorem in [8], the density function is analytic.

For $0 < t < 4$, set $a_1(t) = \Phi_t(z_1(t))$, $a_2(t) = \Phi_t(z_2(t))$. Note that $\eta_{\lambda_t}(\Phi_t(z)) = z$ for $z \in \overline{\Omega_t}$. From (3.32) we see that A_t is the closed arc on $\mathbb{T}$ with endpoints $a_1(t), a_2(t)$ which contains 1. Thus, to prove that $A_{t_1} \subset A_{t_2}$, it is enough to prove that $\arg(a_1(t))$ is an increasing function of t . A direct computation shows that $|z_1(t) - 1|^2 = t$ and $\arg(\Sigma_{\lambda_t}(z_1(t))) = \Im z_1(t) = \sqrt{t(4-t)}/2$. We thus have

$$\arg(a_1(t)) = \Im z_1(t) + \arg(z_1(t)) = \sin(\theta_1(t)) + \theta_1(t).$$

From $z_1(t) = (2 - t + \sqrt{t^2 - 4t})/2$ we see that $\theta_1(t)$ is an increasing function of t . The function $\theta \rightarrow \sin(\theta) + \theta$ is an increasing function on $(0, \pi)$. Thus $\arg(a_1(t))$ is an increasing function of t and (1) is proved.

To prove (2), recall that a probability measure is unimodal if its density with respect to Lebesgue measure has a unique local maximum. η_{λ_t} extends continuously to $\mathbb{T}$, we thus have that

$$(5.4) \quad \frac{d\lambda_t(e^{-i\theta})}{d\theta} = \frac{1}{2\pi} \frac{1 - |\eta_{\lambda_t}(e^{i\theta})|^2}{|1 - \eta_{\lambda_t}(e^{i\theta})|^2}.$$

We first prove the case when $0 < t < 4$. From $\eta_{\lambda_t}(\Phi_t(z)) = z$ for $z \in \overline{\Omega_t}$ and $\eta_{\lambda_t}(1) = x(t)$, to prove λ_t is unimodal, by the boundary correspondence, it is enough to show that the function f_t of u defined by

$$f_t(u) := \frac{1 - |\gamma_t(u)|^2}{|1 - \gamma_t(u)|^2},$$

is a decreasing function on $[0, 1]$ and is maximal at 0. Since $\gamma_t(u) \in \partial\Omega_t$, we have that $|\Phi_t(\gamma_t(u))| = 1$. In other words, we have

$$(5.5) \quad |\gamma_t(u)| \exp\left(\frac{t}{2} f_t(u)\right) = |\gamma_t(u)| \exp\left(\frac{t}{2} \frac{1 - |\gamma_t(u)|^2}{|1 - \gamma_t(u)|^2}\right) = 1.$$

As we shown in Lemma 5.1 that the function $|\gamma_t(u)|$ is an increasing function of u , from (5.5), we deduce that f_t is a decreasing function of u and $\max\{f_t\} = f_t(0)$. By the symmetric property of the function Φ_t in Lemma 5.1, the density function is symmetric with respect to x -axis as well. Thus the density of λ_t has only one local maximum at $\Phi_t(\gamma_t(0)) = \Phi_t(x_1(t)) = 1$.

The proof for the case $t \geq 4$ is similar. In this case $A_t = \mathbb{T}$ and $\max\{f_t\} = f_t(0)$ and $\min\{f_t\} = f_t(1)$. Part (3) is a consequence of Corollary 3.27. $\square$

Remark 7. Biane [17] proved that A_t the support of λ_t is inside an arc of the unit circle, centered at 1, and of width approximately $4\sqrt{t}$ for small value of t .

In fact, the proof of Theorem 5.4 provides a new proof for this fact. From the proof, we see that A_t is the arc on the unit circle which passes through 1 with endpoints

$a_1(t)$ and $a_2(t)$. Also we have $\arg(a_1(t)) = \sin \theta_1(t) + \theta_1(t)$ and $\theta_1(t) = \arg(z_1(t)) \approx \sqrt{t}$ for small value of t . We thus have $|A_t| \approx 4\sqrt{t}$ for small value of t .

5.2. The positive half line case. Let $\lambda \in \mathcal{M}_{\mathbb{R}^+}$ be the analogue of the normal distribution such that $\Sigma_\lambda(z) = \exp\left(\frac{z+1}{z-1}\right)$.

We restate Proposition 6.14 in [16] in terms of η and Σ transforms as follows.

Lemma 5.5. *Let μ be a $\boxtimes$ -infinitely divisible measure on $\mathbb{R}^+$, and set $\Phi_\mu(z) := z\Sigma_\mu(z)$.*

- (1) *We have $\Phi_\mu(\eta_\mu(z)) = z$ for every $z \in \mathbb{C}^+$.*
- (2) *The set $\{\eta_\mu(z) : z \in \mathbb{C}^+\} = \Omega$, where Ω is the component of the set $\{z \in \mathbb{C}^+ : \Im(\Phi_\mu(z)) > 0\}$ whose boundary contains the left half line $(-\infty, 0)$. Moreover, $\eta_\mu(\Phi_\mu(z)) = z$ for $z \in \Omega$.*

We denote $\Phi_\lambda(z) = z \exp\left(\frac{z+1}{z-1}\right)$. The following lemma is elementary.

Lemma 5.6. *Φ_λ has zero of order one at $2 - \sqrt{3}$ and $2 + \sqrt{3}$, and Φ_λ has an essential singularity at 1. These are the only zeros and singularities of Φ_λ .*

Theorem 5.7. *The measure λ has no singular part. The support of this measure is the closure of its interior, and this interior has only one connected component.*

Proof. By Theorem 7.5 in [15], the measure λ has compact support on $\mathbb{R}^+$.

Let Ω be the component of $\{z \in \mathbb{C}^+ : \Im(\Phi_\lambda(z)) > 0\}$ whose boundary contains $(-\infty, 0)$. By Lemma 5.5, $\eta_\lambda : \mathbb{C}^+ \rightarrow \Omega$ is a conformal map and Φ_λ is its inverse map, thus Ω is simply connected. By Lemma 5.6, $\partial\Omega$ is locally analytic. A general theorem in complex analysis tells us that η_λ extends continuously to $\mathbb{C}^+ \cup \mathbb{R}$ and it establishes a homeomorphism between the real axis and $\partial\Omega$. We continue to denote by η_λ and Φ_λ their extensions.

We claim that

$$\partial\Omega = \left(-\infty, 2 - \sqrt{3}\right] \cup \left[2 + \sqrt{3}, +\infty\right) \cup \mathcal{L},$$

where $\mathcal{L}$ is an analytic and open curve in $\mathbb{C}^+$ with endpoints $2 - \sqrt{3}$ and $2 + \sqrt{3}$. We denote $\gamma(t) = \eta_\lambda(t)$, $t \in \mathbb{R}$ be a parametrization of $\partial\Omega$. Set $t_1 = \Phi_\lambda(2 - \sqrt{3}) > 0$ and $t_2 = \Phi_\lambda(2 + \sqrt{3}) > 0$, then $\eta_\lambda(t_1) = 2 - \sqrt{3}$ and $\eta_\lambda(t_2) = 2 + \sqrt{3}$, and $\mathcal{L} = \{\gamma(t)\}_{t_1 < t < t_2}$. Note that

- (1) $(-\infty, 0) \subset \partial\Omega$,
- (2) $\Phi'_\lambda(x) > 0$ for all $x \in (-\infty, 2 - \sqrt{3})$.

From this we deduce that $(-\infty, 2 - \sqrt{3}) \subset \partial\Omega$. Lemma 5.6 tells us Φ_λ has a zero of order one at $2 - \sqrt{3}$, therefore $\partial\Omega$ has a corner of opening $\pi/2$ at $2 - \sqrt{3}$. Note that $\Phi'_\lambda(x) > 0$ for all $x \in (2 + \sqrt{3}, +\infty)$, thus $(2 + \sqrt{3}, +\infty) \subset \partial\Omega$, and $\partial\Omega$ has a corner of opening $\pi/2$ at $2 + \sqrt{3}$.

It remains to prove that $\mathcal{L} \cap \mathbb{R} = \emptyset$. First we show $1 \notin \mathcal{L}$. Suppose that is the case, and suppose $\gamma(t_0) = 1$ where $t_1 < t_0 < t_2$, by continuity, we have

$$\gamma(t) \exp \left(\frac{\gamma(t) + 1}{\gamma(t) - 1} \right) = \Phi_\lambda(\gamma(t)) = \Phi_\lambda(\eta_\lambda(t)) = t$$

for all $t \in \mathbb{R}$. Therefore in a small neighborhood of t_0 , we have

$$\frac{\gamma(t) + 1}{\gamma(t) - 1} = \ln \left(\frac{t}{\gamma(t)} \right).$$

The left hand of the above equation blows up, while the right hand side is bounded. This contradiction tells us that $1 \notin \mathcal{L}$. Now suppose $\mathcal{L}$ touches the real axis at $x_0 \in (2 - \sqrt{3}, 1) \cup (1, 2 + \sqrt{3})$. Since Ω is connected, it is not hard to see that x_0 must be a critical point of Φ_λ . This is not possible by Lemma 5.6. We therefore proved that $\mathcal{L} \subset \mathbb{C}^+$ and the claim.

From the definitions of the Cauchy transform and η -transform, one can easily check that

$$G_\lambda \left(\frac{1}{z} \right) = \frac{z}{1 - \eta_\lambda(z)}.$$

From the above equation we know that G_λ extends to be a continuous function on $\mathbb{C} \cup \mathbb{R}$, and $\{x \in \mathbb{R} : \Im(G_\mu(x)) > 0\} = (1/t_2, 1/t_1)$. By the Stieltjes inverse formula, we deduce that the support of λ is $(1/t_2, 1/t_1)$. From the analyticity of the curve $\mathcal{L} \subset \mathbb{C}^+$, we conclude that λ has positive and analytic density in the interior of its support. $\square$

We are interested in the level curves of the function

$$(5.6) \quad f(r, \theta) = \theta - \frac{2r \sin \theta}{1 - 2r \cos \theta + r^2} = \arg(\Phi_\lambda(z)),$$

where $z = re^{i\theta} \in \mathbb{C}^+$. For $t \leq 0$, set $\gamma_t = \{z = re^{i\theta} \in \mathbb{C}^+ : f(r, \theta) = t\}$.

Proposition 5.8. (A) γ_0 is a simple open curve with endpoints $2 - \sqrt{3}, 2 + \sqrt{3}$ and $\gamma_0 = \mathcal{L}$.

(B) γ_t is a simple open curve which starts at $z = 1$ and ends at $z = 1$ as well for all $t < 0$.

Denote by Ω_0 the open domain bounded $\gamma_0 \cup [2 - \sqrt{3}, 2 + \sqrt{3}]$. For all $t < 0$, Denote by Ω_t the open domain bounded $\gamma_t \cup \{1\}$.

(C) For $t_1 < t_2 \leq 0$, we have that $\Omega_{t_1} \subset \Omega_{t_2}$; and for all $t_0 \leq 0$, $\Omega_{t_0} = \cup_{t < t_0} \Omega_t$.

Proof. Given $\theta \in (0, \pi)$, we define a function of r by $f_\theta(r) = f(r, \theta)$ for $r \in (0, +\infty)$. We first note that $f(r, \theta) < \theta < \pi$ and observe that

$$\lim_{r \rightarrow +\infty} f_\theta(r) = \theta.$$

We thus have that $\{z = re^{i\theta} : f(r, \theta) > 0, 0 < \theta < \pi\} \subset \Phi^{-1}(\mathbb{C}^+)$.

Given $\theta \in (0, \pi)$ and $t \leq 0$, the equation $f(r, \theta) = t$ is equivalent to the quadratic equation

$$(5.7) \quad h_\theta(r) := (\theta - t)r^2 - (2(\theta - t)\cos\theta + 2\sin\theta)r + \theta - t = 0$$

with discriminant $d(\theta, t) = [2(\theta - t)\cos\theta + 2\sin\theta]^2 - 4(\theta - t)^2$. We then rewrite $d(\theta, t)$ as follows.

$$(5.8) \quad d(\theta, t) = 4(1 - \cos^2(\theta)) \left[\frac{\sin\theta}{1 + \cos\theta} + \theta - t \right] \cdot \left[\frac{\sin\theta}{1 - \cos\theta} - \theta + t \right],$$

We observe that the first two factors in (5.8) are never zero for $\theta \in (0, \pi)$, thus only the last factor in (5.8) matters to determine the sign of $d(\theta, t)$. We consider the function k by $k(\theta) = \sin\theta/(1 - \cos\theta) - \theta$ for $\theta \in (0, \pi)$, and calculate

$$(5.9) \quad k'(\theta) = \frac{1}{\cos\theta - 1} - 1 < 0,$$

which implies that k is a decreasing function of θ . For $t \leq 0$, we now set $d_t(\theta) := d(\theta, t)$. We then deduce that $d_t(\theta) = 0$ has exactly one solution, which we denote by θ_t , and $d_t(\theta) > 0$ if and only if $0 < \theta < \theta_t$. Therefore, the half line $r = \theta$ intersects with γ_t at two points if and only if $0 < \theta < \theta_t$ and the half line $r = \theta_t$ is tangent to γ_t . Moreover, $\theta_{t_1} < \theta_{t_2}$ if $t_1 < t_2 \leq 0$.

For the solutions of the equation $f(r, \theta) = 0$, one can check as $\theta \rightarrow 0$, r satisfying the equation $r^2 - 4r + 1$. Given $t < 0$, for the solutions of the equation $f(r, \theta) = t$, we can easily see that r tend to 1 as $\theta \rightarrow 0$. Now (A) and (B) follow from this observation.

Given $\theta \in (0, \pi)$, from (5.6), we see that the function $f_\theta(r)$ defined by $f_\theta(r) = f(r, \theta)$ has exactly one local minimum at $r = 1$. $f_\theta(r)$ is a decreasing function of r on $(0, 1)$ and an increasing function of r on $(1, \infty)$. Therefore, if the half line $r = \theta$ intersects with γ_t at two points, then one of them is inside the unit circle of $\mathbb{C}$ and the other one is outside the unit circle. We conclude that (C) is valid. $\square$

It is interesting to compare the following result with Proposition 2.2 and Lemma 3.2.

Corollary 5.9. *We have that $\Phi_\lambda^{-1}(\mathbb{C}^+) = \Omega \cup_{k=1}^\infty (\Omega_{(2k-1)\pi} \setminus \Omega_{(2k-2)\pi})$. Moreover, Ω and $\Omega_{(2k-1)\pi} \setminus \Omega_{(2k-2)\pi}$ ($k = 1, 2, \dots$) are all connected components of Φ_λ . In particular, $\Phi_\lambda^{-1}(\mathbb{C}^+)$ has infinitely many connected components.*

We would like to point out that for $z = re^{i\theta} \in \mathcal{L} = \gamma_0$, the curve $\mathcal{L}$ can be parametrized by r . Noticing (5.7) and (5.8), we first observe the following equivalence relations:

$$(5.10) \quad d(\theta, 0) = 0 \Leftrightarrow \theta \cos\theta + \sin\theta = \theta \Leftrightarrow r = 1.$$

By (5.9), we see that (5.10) has exactly one solution θ_0 for $\theta \in (0, \pi)$. By differentiating the equation $f(r, \theta) = 0$, we obtain that

$$(5.11) \quad \frac{d\theta}{dr} = \frac{2\theta \cos \theta + 2 \sin \theta - 2\theta r}{r^2 + 2\theta \sin \theta - 4 \cos \theta + 1}.$$

Thus, $d\theta/dr = 0$ if and only if $r = (\theta \cos \theta + \sin \theta)/\theta$. Fix θ , the equation $f_\theta(r) = 0$ is equivalent to the quadratic equation $\theta r^2 - (2\theta \cos \theta + 2 \sin \theta)r + \theta = 0$, from which we deduce that $r = (\theta \cos \theta + \sin \theta)/\theta$ if and only if $d(r, 0) = 0$. From (5.11) and continuity of $d\theta/dr$, we see that $d\theta/dr > 0$ for $0 < \theta < \theta_0, r < 1$ and $d\theta/dr < 0$ for $0 < \theta < \theta_0, r > 1$. Therefore, for the solutions of the equation $f(r, \theta) = 0$, θ is a function of r and the curve $\mathcal{L}$ can be parametrized by r .

We denote by g the density function of λ . From the equation $G_\lambda(1/x) = x/(1 - \eta_\lambda(x))$, we obtain the following formula for the density function of λ .

Proposition 5.10. *Given $z = re^{i\theta} \in \gamma_0 = \mathcal{L}$, we have*

$$g(1/x) = \theta \Phi_\lambda(z) = r\theta \exp \left(\frac{r^2 - 1}{1 - 2r \cos \theta + r^2} \right),$$

where $x = \Phi_\lambda(z)$.





FIGURE 1. Level curves of $g_2(r, \theta) = |\Phi_2(re^{i\theta})|$. The vertical axis indicates θ , and the horizontal axis indicates r .



ACKNOWLEDGEMENTS

I am grateful to my advisor, Professor Hari Bercovici, for encouragement and many helpful discussions which make this work available. I thank Octavio Arizmendi for informing me his paper [5] and Professor Michael Anshelevich for his interest. I would also like to thank Professor Eric Bedford for several helpful conversations and Aimin Huang for his careful reading of a preliminary version of this paper.

REFERENCES

- [1] Michael Anshelevich, *Free evolution on algebras with two states*, J. Reine Angew. Math. **638** (2010), 75–101. MR 2595336 (2012d:46155)
- [2] ———, *Generators of some non-commutative stochastic process*, arXiv:1104.1381v2 math.OA (2011).
- [3] ———, *Two-state free Brownian motions*, J. Funct. Anal. **260** (2011), no. 2, 541–565. MR 2737412 (2012c:46147)
- [4] ———, *Free evolution on algebras with two states II*, arXiv:1204.0289v1 math.OA (2012).
- [5] Octavio Arizmendi, *k-divisible random variables in free probability*, arXiv:1203.4780v1 [math.PR] (2012).
- [6] Octavio Arizmendi and Takahiro Hasebe, *Semigroups related to additive and multiplicative, free and boolean convolutions*, arXiv:1105.3344v3 [math.PR] (2012).
- [7] Serban T. Belinschi and Hari Bercovici, *Atoms and regularity for measures in a partially defined free convolution semigroup*, Math. Z. **248** (2004), no. 4, 665–674. MR 2103535 (2006i:46095)
- [8] ———, *Partially defined semigroups relative to multiplicative free convolution*, Int. Math. Res. Not. (2005), no. 2, 65–101. MR 2128863 (2006f:46061)
- [9] ———, *A new approach to subordination results in free probability*, J. Anal. Math. **101** (2007), 357–365. MR 2346550 (2008i:46059)
- [10] Serban T. Belinschi and Alexandru Nica, *η -series and a Boolean Bercovici-Pata bijection for bounded k-tuples*, Adv. Math. **217** (2008), no. 1, 1–41. MR 2357321 (2009c:46088)
- [11] ———, *On a remarkable semigroup of homomorphisms with respect to free multiplicative convolution*, Indiana Univ. Math. J. **57** (2008), no. 4, 1679–1713. MR 2440877 (2009f:46087)
- [12] ———, *Free Brownian motion and evolution towards $\boxplus$ -infinite divisibility for k-tuples*, Internat. J. Math. **20** (2009), no. 3, 309–338. MR 2500073 (2010g:46108)
- [13] Hari Bercovici, *On Boolean convolutions*, Operator theory 20, Theta Ser. Adv. Math., vol. 6, Theta, Bucharest, 2006, pp. 7–13. MR 2276927 (2007m:46105)
- [14] Hari Bercovici and Vittorino Pata, *Stable laws and domains of attraction in free probability theory*, Ann. of Math. (2) **149** (1999), no. 3, 1023–1060, With an appendix by Philippe Biane. MR 1709310 (2000i:46061)
- [15] Hari Bercovici and Dan Voiculescu, *Lévy-Hinčin type theorems for multiplicative and additive free convolution*, Pacific J. Math. **153** (1992), no. 2, 217–248. MR 1151559 (93k:46052)
- [16] ———, *Free convolution of measures with unbounded support*, Indiana Univ. Math. J. **42** (1993), no. 3, 733–773. MR 1254116 (95c:46109)
- [17] Philippe Biane, *Free Brownian motion, free stochastic calculus and random matrices*, **12** (1997), 1–19. MR 1426833 (97m:46104)
- [18] ———, *On the free convolution with a semi-circular distribution*, Indiana Univ. Math. J. **46** (1997), no. 3, 705–718. MR 1488333 (99e:46084)
- [19] ———, *Processes with free increments*, Math. Z. **227** (1998), no. 1, 143–174. MR 1605393 (99e:46085)

- [20] Gennadii P. Chistyakov and Friedrich Gtze, *The arithmetic of distributions in free probability theory*, Central European Journal of Mathematics **9** (2011), 997–1050, 10.2478/s11533-011-0049-4.
- [21] ———, *Asymptotic expansion in the CLT in free probability*, arXiv:1109.4844v2 [math.PR] (2011).
- [22] Uwe Franz, *Boolean convolution of probability measures on the unit circle*, Analyse et probabilités, Sémin. Congr., vol. 16, Soc. Math. France, Paris, 2008, pp. 83–94. MR 2599263 (2011c:46136)
- [23] Einar Hille, *Analytic function theory. Vol. 1*, Introduction to Higher Mathematics, Ginn and Company, Boston, 1959. MR 0107692 (21 #6415)
- [24] Romuald Lenczewski, *Decompositions of the free additive convolution*, J. Funct. Anal. **246** (2007), no. 2, 330–365. MR 2321046 (2008d:28009)
- [25] Hans Maassen, *Addition of freely independent random variables*, J. Funct. Anal. **106** (1992), no. 2, 409–438. MR 1165862 (94g:46069)
- [26] Alexandru Nica, *Multi-variable subordination distributions for free additive convolution*, J. Funct. Anal. **257** (2009), no. 2, 428–463. MR 2527024 (2010j:46121)
- [27] Alexandru Nica and Roland Speicher, *On the multiplication of free N -tuples of noncommutative random variables*, Amer. J. Math. **118** (1996), no. 4, 799–837. MR 1400060 (98i:46069)
- [28] N. Raj Rao and Roland Speicher, *Multiplication of free random variables and the S -transform: the case of vanishing mean*, Electron. Comm. Probab. **12** (2007), 248–258. MR 2335895 (2008f:46082)
- [29] Roland Speicher and Reza Woroudi, *Boolean convolution*, Free probability theory (Waterloo, ON, 1995), Fields Inst. Commun., vol. 12, Amer. Math. Soc., Providence, RI, 1997, pp. 267–279. MR 1426845 (98b:46084)
- [30] Dan Voiculescu, *Addition of certain noncommuting random variables*, J. Funct. Anal. **66** (1986), no. 3, 323–346. MR 839105 (87j:46122)
- [31] Jiun-Chau Wang, *Limit laws for Boolean convolutions*, Pacific J. Math. **237** (2008), no. 2, 349–371. MR 2421126 (2009h:46128)
- [32] ———, *Local limit theorems in free probability theory*, Ann. Probab. **38** (2010), no. 4, 1492–1506. MR 2663634 (2011i:46081)

DEPARTMENT OF MATHEMATICS, RAWLES HALL, 831 EAST THIRD STREET, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47405, U.S.A.

E-mail address: pzhong@indiana.edu